



Northeast Fisheries Science Center Technical Memorandum 350

Management Track Assessments Fall 2022

by Northeast Fisheries Science Center

September 2022

Management Track Assessments Fall 2022

by Northeast Fisheries Science Center

U.S. Department of Commerce
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Northeast Fisheries Science Center
Woods Hole, Massachusetts

September 2022

NOAA Technical Memorandum, Editorial Notes

Editorial Treatment: In the interest of expedited publication, this report has undergone a truncated version of the NEFSC Editorial Office's typical technical and copy editing procedure. Aside from the front and back matter included in this document, all writing and editing have been performed by the authors included on the title page.

Information Quality Act Compliance: In accordance with section 515 of Public Law 106-554, the NEFSC completed both technical and policy reviews for this report. These predissemination reviews are on file at the NEFSC Editorial Office.

Species Names: The NEFSC Editorial Office's policy on the use of species names in all technical communications is generally to follow the American Fisheries Society's lists of scientific and common names for fishes, mollusks, and decapod crustaceans and to follow the Society for Marine Mammalogy's guidance on scientific and common names for marine mammals. Exceptions to this policy occur when there are subsequent compelling revisions in the classifications of species, resulting in changes in the names of species.

Statistical Terms: The NEFSC Editorial Office's policy on the use of statistical terms in all technical communications is generally to follow the International Standards Organization's handbook of statistical methods.

This document may be cited as:

NEFSC. 2022. Management Track Assessments Fall 2022. US Dept Commer, Northeast Fish Sci Cent Tech Memo. 350; 167p.+xv. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at <http://www.nefsc.noaa.gov/nefsc/publications/>.

TABLE OF CONTENTS

Page

LISTS

List of Tables.	iii
List of Figures	v

GLOSSARIES

Abbreviations and Acronyms	viii
Abbreviations for fish stocks reviewed	xi
Statistical/review concepts, parameters, etc.	xii
Locations/regions: state, country, etc.	xv

DOCUMENT CONTENTS

1. Panel Report.	1
1.1. Executive Summary	1
Appendix A. AOP Meetings Summary	4
<i>Appendix A.1. Meeting participants</i>	14
Appendix B. Terms of Reference.	15
Appendix C. Peer Review Meeting Attendees.	16
Appendix D. Peer Review Meeting Agenda	18
2. Atlantic wolffish	19
2.1. Reviewer Comments: Atlantic wolffish	22
3. Gulf of Maine winter flounder	28
3.1. Reviewer Comments: Gulf of Maine winter flounder	31
4. Georges Bank Winter Flounder	37
4.1. Reviewer Comments: Georges Bank Winter Flounder	41
5. Georges Bank haddock	48
5.1. Reviewer Comments: Georges Bank haddock	52
6. Gulf of Maine haddock	59
6.1. Reviewer Comments: Gulf of Maine haddock.	62
7. Atlantic halibut	71
7.1. Reviewer Comments: Atlantic halibut	74
8. White hake	80
8.1. Reviewer Comments: White hake	83
9. Northern Monkfish	92

9.1. Reviewer Comments: Northern Monkfish	94
10. Southern Monkfish	98
10.1. Reviewer Comments: Southern Monkfish	100
11. Ocean pout	104
11.1. Reviewer Comments: Ocean pout	106
12. American plaice	111
12.1. Reviewer Comments: American plaice	114
13. Pollock	121
13.1. Reviewer Comments: Pollock	125
14. Witch flounder	134
14.1. Reviewer Comments: Witch flounder	136
15. Cape Cod–Gulf of Maine yellowtail flounder	142
15.1. Reviewer Comments: Cape Cod–Gulf of Maine yellowtail flounder	145
16. Southern New England/Mid-Atlantic yellowtail flounder	152
16.1. Reviewer Comments: Southern New England/Mid-Atlantic yellowtail flounder	156
Photo Gallery	164



Healthy seafood delights: shrimp, mussels, scallop, and fish dish.

LIST OF TABLES

1	Stocks reviewed at September 2022 Management Track Assessment Peer Review meeting	3
2	Stocks reviewed at September 2022 Management Track Assessment Stock Assessments meeting	5
3	Catch and status table for Atlantic wolffish	19
4	Estimated reference points for Atlantic wolffish	19
5	Catch and status table for Gulf of Maine winter flounder	28
6	Estimated reference points for Gulf of Maine winter flounder	28
7	Catch and status table for Georges Bank Winter Flounder	37
8	Estimated reference points for Georges Bank Winter Flounder	38
9	Short term projections for Georges Bank Winter Flounder	38
10	Catch and status table for Georges Bank haddock	48
11	Estimated reference points for Georges Bank haddock	49
12	Short term projections for Georges Bank haddock	49
13	Catch and status table for Gulf of Maine haddock	59
14	Estimated reference points for Gulf of Maine haddock	60
15	Short term projections for Gulf of Maine haddock	60
16	Catch and status table for Atlantic halibut	71
17	Estimated reference points for Atlantic halibut	71
18	Catch and status table for white hake	80
19	Estimated reference points for white hake	81
20	Short term projections for white hake	81
21	Catch and status table for Northern Monkfish	92
22	Estimated reference points for Northern Monkfish	92
23	Catch and status table for Southern Monkfish	98
24	Estimated reference points for Southern Monkfish	98
25	Catch and status table for ocean pout	104
26	Estimated reference points for ocean pout	104
27	Catch and status table for American plaice	111

28 Estimated reference points for American plaice 112

29 Short term projections for American plaice 112

30 Catch and status table for pollock 121

31 Estimated reference points for pollock 122

32 Short term projections for pollock 122

33 Catch and status table for witch flounder 134

34 Estimated reference points for witch flounder 134

35 Catch and status table for Cape Cod–Gulf of Maine yellowtail flounder 142

36 Estimated reference points for Cape Cod–Gulf of Maine yellowtail flounder 143

37 Short term projections for Cape Cod–Gulf of Maine yellowtail flounder 143

38 Catch and status table for Southern New England-Mid Atlantic yellowtail flounder 152

39 Estimated reference points for Southern New England/Mid-Atlantic yellowtail flounder . . . 153

40 Short term projections for Southern New England/Mid-Atlantic yellowtail flounder 153



The reason behind it all.

LIST OF FIGURES

1	Estimated trends in biomass for Atlantic wolffish	23
2	Estimated trends in fishing mortality for Atlantic wolffish	24
3	Trends in estimated recruitment for Atlantic wolffish	25
4	Total catch of Atlantic wolffish	26
5	Indices of abundance for Atlantic wolffish	27
6	Estimated trends in biomass for Gulf of Maine winter flounder	33
7	Estimated trends in fishing mortality for Gulf of Maine winter flounder	34
8	Total catch of Gulf of Maine winter flounder	35
9	Indices of abundance for Gulf of Maine winter flounder	36
10	Estimated trends in biomass for Georges Bank Winter Flounder	43
11	Estimated trends in fishing mortality for Georges Bank Winter Flounder	44
12	Trends in estimated recruitment for Georges Bank Winter Flounder	45
13	Total catch of Georges Bank Winter Flounder	46
14	Indices of abundance for Georges Bank Winter Flounder	47
15	Estimated trends in biomass for Georges Bank haddock	54
16	Estimated trends in fishing mortality for Georges Bank haddock	55
17	Trends in estimated recruitment for Georges Bank haddock	56
18	Total catch of Georges Bank haddock	57
19	Indices of abundance for Georges Bank haddock	58
20	Estimated trends in biomass for Gulf of Maine haddock	66
21	Estimated trends in fishing mortality for Gulf of Maine haddock	67
22	Trends in estimated recruitment for Gulf of Maine haddock	68
23	Total catch of Gulf of Maine haddock	69
24	Indices of abundance for Gulf of Maine haddock	70
25	Estimated trends in biomass for Atlantic halibut	76
26	Estimated trends in fishing mortality for Atlantic halibut	77
27	Total catch of Atlantic halibut	78
28	Indices of abundance for Atlantic halibut	79

29	Estimated trends in biomass for white hake	87
30	Estimated trends in fishing mortality for white hake	88
31	Trends in estimated recruitment for white hake	89
32	Total catch of white hake	90
33	Indices of abundance for white hake	91
34	Total catch of Northern Monkfish	96
35	Indices of abundance for Northern Monkfish	97
36	Total catch of Southern Monkfish	102
37	Indices of abundance for Southern Monkfish	103
38	Estimated trends in biomass for ocean pout	107
39	Estimated trends in fishing mortality for ocean pout	108
40	Total catch of ocean pout	109
41	Indices of abundance for ocean pout	110
42	Estimated trends in biomass for American plaice	116
43	Estimated trends in fishing mortality for American plaice	117
44	Trends in estimated recruitment for American plaice	118
45	Total catch of American plaice	119
46	Indices of abundance for American plaice	120
47	Estimated trends in biomass for pollock	129
48	Estimated trends in fishing mortality for pollock	130
49	Trends in estimated recruitment for pollock	131
50	Total catch of pollock	132
51	Indices of abundance for pollock	133
52	Estimated trends in biomass for witch flounder	138
53	Estimated trends in fishing mortality for witch flounder	139
54	Total catch of witch flounder	140
55	Indices of abundance for witch flounder	141
56	Estimated trends in biomass for Cape Cod–Gulf of Maine yellowtail flounder	147
57	Estimated trends in fishing mortality for Cape Cod–Gulf of Maine yellowtail flounder	148
58	Trends in estimated recruitment for Cape Cod–Gulf of Maine yellowtail flounder	149

59 Total catch of Cape Cod–Gulf of Maine yellowtail flounder 150

60 Indices of abundance for Cape Cod–Gulf of Maine yellowtail flounder 151

61 Estimated trends in biomass for Southern New England/Mid-Atlantic yellowtail flounder . 159

62 Estimated trends in fishing mortality for Southern New England/Mid-Atlantic yellowtail flounder 160

63 Trends in estimated recruitment for Southern New England/Mid-Atlantic yellowtail flounder 161

64 Total catch of Southern New England/Mid-Atlantic yellowtail flounder 162

65 Indices of abundance for Southern New England/Mid-Atlantic yellowtail flounder 163



NOAA research vessel Henry B. Bigelow underway

Abbreviations and Acronyms

- Albatross** refers to activities of the **NOAA** vessel **Albatross IV** 2, 51, 84, 85, 127
- Albatross IV** Research vessel **NOAAS** Albatross IV, in service until November 2008 viii, 125
- AOP** Assessment Oversight Panel 4–14, 20, 31, 32, 41, 53, 62, 74, 83, 95, 101, 105, 106, 113, 115, 124, 125, 156
- ASMFC** Atlantic States Marine Fisheries Commission 2, 5, 12, 14, 15, 82, 83, 91
- Bigelow** refers to activities of the **NOAA** vessel **Henry B. Bigelow** 2, 10, 20, 61, 84, 85, 105, 125, 127
- BTS** bottom trawl survey 2, 84, 94, 100
- CAMS** Catch Accounting and Monitoring System 2, 5–7, 9–13, 20, 30, 39, 83, 105, 113, 144
- CIE** Center for Independent Experts 13, 61
- CJFAS** Canadian Journal of Fisheries and Aquatic Sciences 94, 100
- CRD** Center Reference Document **CLXXXV**
- CSE** Council of Science Editors **CLXXXV**
- DFO** Department of Fisheries and Oceans, Canadian 7, 8, 17, 38, 39, 47, 58, 74, 146
- FSD** Fisheries Statistics Division 8, 71–74, 76, 77
- F/V** fishing vessel 127
- GARFO** Greater Atlantic Regional Fisheries Office 1, 14, 15, 17
- GARM** Groundfish Assessment Review Meeting 6, 31
- GARM III** 3rd Groundfish Assessment Review Meeting, 2008 10, 52, 53, 106, 144–146, 157
- H.B. Bigelow** refers to activities of the **NOAA** vessel **Henry B. Bigelow** 51
- Henry B. Bigelow** **NOAA** research vessel Henry B. Bigelow, with specialized trawling net mechanisms; commissioned July 2007, used for surveys 2009–2019 vii, viii, 164
- ICES** International Council for the Exploration of the Sea (European Union) 157
- MA DMF** Massachusetts Division of Marine Fisheries 6, 7, 10, 14–17, 27, 28, 31, 33, 34, 36, 143, 144, 151
- MAFMC** Mid-Atlantic Fisheries Management Council 4, 14, 16
- ME DMR** Maine Department of Marine Resources 14, 16, 17
- MRFSS** Marine Recreational Fisheries Statistics Survey 12
- MRIP** Marine Recreational Information Program 7, 123

MT Management Track 63, 64

NAFO Northwest Atlantic Fisheries Organization xii

NCDMF North Carolina Division of Marine Fisheries 17

NDPSWG Northeast Data Poor Stocks Working Group 6, 19, 20, 22

NEFMC New England Fisheries Management Council 1, 4, 14–17, 106, 111, 135

NEFOP Northeast Fishery Observer Program 2, 6

NEFSC Northeast Fisheries Science Center III, 1, 2, 5–12, 14–17, 19–22, 27–31, 33, 34, 36, 37, 39, 41, 42, 47, 48, 50–52, 58, 59, 61–63, 70, 79, 80, 83, 85, 91, 94, 97, 100, 103–106, 110, 111, 113, 114, 120, 121, 125, 126, 133–136, 141, 143–146, 151, 152, 154, 156, 163, CLXXXIV, CLXXXV

NMFS National Marine Fisheries Service 17, 64, 74, 85, 92, 94, 95, 98, 100, 101, 164, CLXXXIV

NOAA National Oceanographic and Atmospheric Administration vii–ix, xi, 1, 2, 4, 16, 41, 52, 53, 62, 71, 83, 86, 106, 137, 146, 156, 157, 164–CLXXXIV

NOAAS NOAA ship viii

NRCC Northeast Regional Coordinating Council 4

PDT Plan Development Team 12, 135

RT Research Track 63

R/V research vessel 61

SARC 50 50th Stock Assessment Review Committee meeting, 2010 121

SARC 52 52nd Stock Assessment Review Committee meeting, 2011 28, 29, 31

SASINF Stock Assessment Support Information 1, 114

SAW 50 50th Stock Assessment Workshop, 2010 5, 126, 128

SAW 52 52nd Stock Assessment Workshop, 2011 32

SAW 54 54th Stock Assessment Workshop, 2012 11, 156, 157

SAW 56 56th Stock Assessment Workshop, 2013 11, 12, 83, 86

SAW 66 66th Stock Assessment Workshop, 2019 146

SMAST School for Marine Science and Technology (New Bedford, Maine) 15, 17

SSC Scientific and Statistical Committee 14, 15, 64, 85, 106, 127, 135

TMGC Transboundary Management Guidance Committee 16

TOR Term of Reference 9, 62–64, 83–85, 125–127

TRAC Transboundary Resources Assessment Committee 146

WHAM Woods Hole Assessment Model 5, 9, 13, 48, 49, 52, 53, 111, 113, 114, 155, 157

WHOI Woods Hole Oceanographic Institute, MA x, 17, 164



Aerial view of Woods Hole Oceanographic Institute, MA; photo ©WHOI

Abbreviations for fish stocks reviewed

These are the abbreviations for fish stock names, as seen in the footers of each of the fish stock reports.

CATUNIT (*Anarhichas lupus*) Atlantic wolffish 19–27

FLWGB (*Pseudopleuronectes americanus*) winter flounder, from the Georges Bank 37–47

FLWGM (*Pseudopleuronectes americanus*) winter flounder, from the Gulf of Maine 28–36

HADGB (*Melanogrammus aeglefinus*) haddock, from the Georges Bank 48–58

HADGM (*Melanogrammus aeglefinus*) haddock, from the Gulf of Maine 59–70

HALUNIT (*Hippoglossus hippoglossus*) Atlantic halibut 71–79

HKWUNIT (*Urophycis tenuis*) white hake 80–91

MNKN (*Lophius americanus*) Northern monkfish 92–97

MNKS (*Lophius americanus*) Southern monkfish 98–103

OPTUNIT (*Zoarces americanus*) ocean pout 104–110

PLAUNIT (*Hippoglossoides platessoides*) American plaice 111–120

POKUNIT (*Pollachius virens*) pollock 121–133

WITUNIT (*Glyptocephalus cynoglossus*) witch flounder 134–141

YELCCGM (*Limanda ferruginea*) yellowtail flounder, from Cape Cod to Gulf of Maine 142–151


YELSNEMA (*Limanda ferruginea*) yellowtail flounder, from Southern New England to Mid-Atlantic 152–163



Atlantic Wolffish



Winter Flounder




Haddock



Atlantic Halibut



White Hake



Monkfish



Ocean Pout



American Plaice



Atlantic Pollock



Witch Flounder



Yellowtail Flounder

Images from [NOAA Fisheries](#) and [FishWatch.gov](#).

Statistical/review concepts, parameters, etc.

- 000s thousands 37, 38, 45, 48, 49, 56, 59, 60, 68, 80, 81, 89, 111, 112, 118, 121, 122, 131, 142, 143, 149, 152, 153, 161
- 5Z NAFO subdivision 5Z, having subareas 5Ze, 5Zej, etc. 73, 74
- AA Annual Allocation 7, 10–12, 113
- ABC acceptable biological catch 7, 29, 94, 100
- ACL annual catch limit 125
- adapt sum-of-squares approach to fitting VPA models 37, 41
- agepro Age-Structured Projection Model, software module 5, 127
- ALK age-length-key 11, 12, 83, 144
- A/L age to length criterion 38, 39
- ASAP Age-Structured Assessment Program, modelling software 5, 9–13, 59, 62, 63, 80–85, 121, 122, 125–127, 152, 155, 156
- ASM At Sea Monitoring 2
- BLLS Bottom Longline Survey 61, 63, 64, 83, 84
- B_{MSY} biomass maximum sustainable yield 11, 28, 104
- BRP biological reference point 5, 10, 15, 63, 84, 106, 127, 146
- BSIA Best Scientific Information Available 32, 41, 53, 62, 74, 83, 95, 101, 115, 125, 156
- $B_{Terminal}$ terminal year biomass 20
- CAA Catches-at-age 5, 12, 82–85, 126
- CDF cumulative distribution function 38
- CI confidence interval 49
- + cm catch at least of specified length in centimeters 7, 28, 29, 31, 33
- Covid refers to coronavirus pandemic years, 2020–2021 2, 5–7, 9, 10, 12, 29, 39, 40, 72, 135, 143
- CPUE catch per unit effort 74
- CV coefficient of variation 62, 82, 85, 126
- E_{Full} exploitation mortality on fully selected ages 28, 34
- $E_{MSY proxy}$ the exploitation rate commensurate with fishing at the proxy for maximum sustainable yield 28, 29, 31, 34

$E_{40\%}$ exploitation rate at 40% of the total catch 28, 29, 31

F (instantaneous) fishing mortality rate 20, 29, 37–39, 41, 48–50, 52, 60, 61, 63, 67, 72, 81, 82, 92, 98, 105, 112, 113, 121–123, 126, 127, 130, 135, 143, 144, 152–154

F_{AVG} averaged fishing mortality 121–123, 130

$\bar{F}_{5:7}$ average fishing mortality for fish aged 5 to 7 years 48–50, 55

F_{Full} fishing mortality rate on fully selected ages 19, 20, 24, 29, 37–39, 44, 59, 60, 80–82, 88, 93, 99, 105, 111, 112, 117, 135, 142–144, 148, 152–154, 160

flat sel flat-topped survey selectivity 121–124, 126, 127, 129–131

F_{MSY} fishing mortality rate for maximum sustainable yield 7, 20, 38, 64, 74, 94, 100, 104, 106, 122, 127

$F_{MSY\ proxy}$ proxy estimate of fishing mortality rate for maximum sustainable yield 19, 24, 37–39, 41, 44, 48, 49, 52, 55, 59, 60, 63, 67, 71, 80, 81, 84, 88, 92, 98, 104, 108, 111, 112, 114, 117, 121, 122, 130, 134, 142, 143, 148, 152, 153, 156, 160

F_{ρ} rho-adjusted value for the fishing rate 39, 50, 113, 123, 144, 154

F_{Target} theoretically ideal fishing mortality level for sustainability 44

$F_{Threshold}$ threshold fishing mortality level that indicates overfishing status 24, 34, 44, 55, 67, 88, 108, 117, 130, 148, 160

$F_{40\%SPR}$ fishing mortality for 40% of spawning potential rate 5, 19, 38, 49, 51, 52, 59, 60, 63, 64, 81, 112, 121, 122, 127, 143, 153

GMRF Gaussian Markov random field 49–53

lsmooth renaming of **PlanBsmooth**: a model using log-linear regression and **Loess** smoothing 92–95, 98–100

kg/tow kilograms per tow 58, 104, 110

Loess loess curve fitting (local polynomial regression) xiii, xiv, 6, 12, 94, 100

log-normal probability distribution whose logarithm is normally distributed 21, 27, 36, 58, 66–68, 70, 87–89, 91, 97, 103, 110, 116, 118, 120, 129–131, 133, 141, 159–161, 163

M (instantaneous) natural mortality rate 11, 63, 64

MCMC Markov Chain Monte Carlo analysis 20, 21

ρ Mohn’s rho parameter: the average relative bias of retrospective estimates 20, 39, 48, 50, 60, 63, 66, 67, 80, 82, 85, 113, 123, 144, 154, 155

MSY maximum sustainable yield 19, 28, 38, 49, 60, 71, 81, 92, 98, 104, 112, 122, 134, 143, 153

mt metric ton 19, 28, 29, 31, 37, 38, 41, 43, 46, 48, 49, 52, 59, 60, 63, 64, 71, 74, 80, 81, 84, 85, 92, 98, 104, 111, 112, 114, 121–123, 125, 127, 134–137, 142, 143, 152–154, 156

NA not applicable 28, 71, 92, 98, 134

NAA Numbers-at-age 62

OFL overfishing limit 7, 20, 29, 31, 81, 106

PlanBsmooth ‘Plan B’ model using log-linear regression and **Loess** smoothing xiii, 9, 94, 100

q catchability coefficient 28–32, 136, 155

R expected recruitment numbers 11

R_{MSY} expected recruitment when biomass is consistent with maximum sustainable yield 49

SBRM Standardized Bycatch Reporting Methodology 5, 83

scale Statistical Catch at Length Model, software module 6, 19–21

scall dr scallop dredge 37

SSB spawning stock biomass 5, 19, 20, 29, 37–39, 41, 48–52, 59–61, 63, 64, 66, 71, 80–82, 84, 85, 93, 99, 105, 111–113, 121–123, 126, 127, 135, 142–145, 152–155

SSB_{MSY} spawning stock biomass consistent with maximum sustainable yield 19, 49, 60, 71, 74, 81, 92, 94, 98, 100, 104, 112, 122, 127, 134, 143, 153, 156

$SSB_{MSY proxy}$ proxy value for spawning stock biomass estimation for maximum sustainable yield 19, 23, 37, 38, 41, 43, 48, 52, 54, 59, 60, 63, 66, 80, 84, 87, 104, 106, 111, 114, 116, 121, 129, 142, 143, 146, 152, 159

$SSB/SSB_{Threshold}$ ratio of spawning stock biomass to spawning stock biomass threshold 5

SSB_{ρ} spawning stock biomass level adjusted according to Mohn’s rho 39, 50, 82, 113, 123, 144, 154

SSB_{Target} theoretically ideal spawning stock biomass level 23, 43, 54, 66, 87, 106, 116, 129, 145, 146, 159

$SSB_{Threshold}$ threshold for spawning stock biomass that indicates overfished status 23, 43, 54, 66, 87, 106, 116, 129, 145, 146, 159

VPA virtual population analysis xii, 5, 7, 9, 10, 37, 38, 41, 49, 142–144

WAA Weight-at-age 52, 53, 61–64, 83, 85, 114, 156

Y_c catch years 37

Locations/regions: state, country, etc.

CA Canada xv, 7, 37, 40, 71

CCGM Cape Cod to Gulf of Maine 5, 10, 145

CT Connecticut xv

GB Georges Bank xv, 38

GOM Gulf of Maine xv, 18, 63

MA Massachusetts III, x, xv, 17, 164

MAB Mid-Atlantic Bight xv

ME Maine xv

ME/NH Maine and New Hampshire 7, 10, 28, 31, 33, 34, 36, 85, 143

NH New Hampshire xv

NJ New Jersey xv

NY New York xv

RI Rhode Island xv, 15, 17


SNE Southern New England xv

SNEMA Southern New England and Mid-Atlantic Bight
2, 3, 5, 11, 156

US United States 7, 8, 31, 37–39, 41, 42, 48, 51, 52, 57, 62, 72, 74, 83, 109, 126, 156, 157, 162

VA Virginia 1

VT Vermont xv



1. 2022 MANAGEMENT TRACK PEER REVIEW PANEL REPORT

Richard Merrick¹ (chair), Matt Cieri², Yan Jiao³ and Cate O’Keefe⁴.

1.1. Executive Summary

Eleven fish stock assessments were reviewed by the September 2022 Management Track peer review panel. Eight of these were Level 2 Expedited Reviews: Gulf of Maine and Georges Bank winter flounder (*Pseudopleuronectes americanus*), Atlantic halibut (*Hippoglossus hippoglossus*), Georges Bank haddock (*Melanogrammus aeglefinus*), north and south monkfish (*Lophius piscatorius*), Southern New England/Mid-Atlantic yellowtail flounder (*Limanda ferruginea*), and American plaice (*Hippoglossoides platessoides*). The remaining three stocks received Level 3 Enhanced Review: white hake (*Urophycis tenuis*), Gulf of Maine haddock (*Melanogrammus aeglefinus*), and pollock (*Pollachius virens*). Levels of review were as recommended by the Assessment Oversight Panel [Appendix A](#).

The Peer Review Panel (Panel) for the September 2022 Management Track Assessments met via webinar on September 19–22, 2022. The Panel was to determine whether the completed management track assessment was technically sufficient to (a) evaluate stock status, (b) provide scientific advice and (c) successfully address the assessment Terms of Reference [Appendix B](#). Tables 1 and 2 present a list of the stocks, names of the lead analyst/presenters, and conclusions about stock status and the assessment.

Attendance at the meeting is provided in [Appendix C](#) with the Agenda shown in [Appendix D](#).

We thank Russ Brown (Population Dynamics Branch Chief) and Michele Traver (Assessment Process Lead) for their support during the meeting and to the staff of the Population Dynamics Branch at [NEFSC](#) for the open and collaborative spirit with which they engaged the Panel. Dr. Brown’s presentation on Data Changes was especially appreciated.

Our thanks also extend to the rapporteurs for taking extensive notes during the meeting and to staff of the New England Fishery Management Council ([NEFMC](#)) or [NOAA](#) Fisheries Greater Atlantic Regional Fisheries Office ([GARFO](#)) who provided context and additional background.

The Panel has suggestions for improvements that should be made for future Management Track Assessments with respect to information needs:

1. The [SASINF](#) portal is an incredible asset for these reviews, and we support its continued maintenance. It is not unusual for documents and data to change on the drive during the period of the review, and as such, it would be useful if a version control mechanism was implemented to allow the reviewers to be notified when changes are made to documents on the site.

¹NOAA Fisheries Service (retired)

²Maine Department of Natural Resources

³Dept. of Fish and Wildlife Conservation, Virginia Polytechnic Institute and State University, Blacksburg, [VA](#)

⁴Fishery Applications Consulting Team

2. For transboundary stocks, it would be useful to have a presentation of the science and management for the Canadian fishery.
3. For species with multiple stocks, consider providing an overview of stock status, structure, etc. at the beginning of the stocks' presentations.

The Panel also has several cross-cutting recommendations with respect to the individual stock assessments:

1. Assessment analysts should consider splitting the bottom trawl time series into two stanzas, namely **Albatross** versus **Bigelow** for those stocks where calibration between the two vessels surveys results was weak (e.g., pollock and white hake).
2. The **NEFSC** Bottom Longline Survey should be continued and considered for incorporation in future stock specific Management Track assessments once the time-series has grown.
3. The **ASMFC** shrimp survey provides valuable information on early year-classes for several species and should continue to be supported by **NOAA** (and perhaps renamed to the 'Summer Survey').
4. Reduction in Port sampling for individual lengths and age structures represents a significant threat to the stock assessment enterprise. **NOAA** should decide whether it can return Port sampling to levels comparable with those achieved prior to 2019. If they cannot, they should increase catch sampling by observers (either **ASM** or **NEFOP**) to balance the loss of these data.
5. **NOAA** should continue to evaluate the use of dynamic reference points with analytic assessments.
6. Assessments for stocks at very low abundance with low fishery mortality rates, showed sharp increases in abundance in projection years (e.g., Gulf of Maine winter flounder, **SNEMA** yellowtail flounder). This is a highly uncertain prediction because these increases may be an artifact of the model considering that low fishing mortality directly leads to increased abundance.

The Panel considered general data changes that were applied across assessments, including:

1. adaptation to survey indices resulting from the missing 2020 research surveys due to the **Covid-19** pandemic;
2. increased uncertainty in catch related indices resulting from reduced Port, **NEFOP** or **ASM** observer, and recreational intercept sampling in 2020;
3. use of the Catch Accounting and Monitoring System (**CAMS**) data for commercial landings for 2020 and 2021; and
4. revised swept-area adjusted survey indices for the **NEFSC Bigelow** Bottom Trawl Surveys (**BTS**).

Table 1: Stocks reviewed at September 2022 Management Track Assessment Peer Review meeting

Stock	Lead Analyst/Presenter	Peer review conclusion
Gulf of Maine winter flounder Expedited Review	Paul Nitschke	Stock's overfished status is unknown but overfishing is not occurring
Georges Bank winter flounder Expedited Review	Alex Hansell	Stock is not overfished and overfishing is not occurring
Atlantic halibut Expedited Review	Dan Hennen	Stock status is unknown
Georges Bank haddock Expedited Review	Liz Brooks	Stock is not overfished and overfishing is not occurring
Monkfish – North Expedited Review	Jon Deroba	Stock status is unknown
Monkfish – South Expedited Review	Jon Deroba	Stock status is unknown
SNEMA yellowtail flounder Expedited Review	Chris Legault	Stock is overfished but overfishing is not occurring
American plaice	Larry Alade	Stock is not overfished and overfishing is not occurring
White hake Enhanced Review	Kathy Sosebee	Stock is not overfished and overfishing is not occurring
Gulf of Maine haddock Enhanced Review	Charles Perretti	Stock is not overfished and overfishing is occurring
Pollock Enhanced Review	Brian Linton	Stock is not overfished and overfishing is not occurring

Appendix A. Summary of Assessment Oversight Panel Meetings for September 2022 Management Track Stock Assessments

The **NRCC** Assessment Oversight Panel (**AOP**) met to review the operational stock assessment plans for ocean pout, Atlantic wolffish, Georges Bank winter flounder, Gulf of Maine winter flounder, Cape Cod/Gulf of Maine yellowtail flounder, Southern New England/Mid-Atlantic yellowtail flounder, northern and southern monkfish, Georges Bank haddock, Gulf of Maine haddock, Atlantic halibut, witch flounder, white hake and pollock stocks on May 23–24, 2022. The **AOP** also met on August 3, 2022 to review the assessment plan for American Plaice, which underwent a Research Track peer review in July 2022. Four assessments were recommended for Level 1 Reviews (Direct Delivery) and these assessments will undergo an internal review before being delivered to the appropriate management body. The assessments for stocks/species recommended for Level 2 and 3 peer reviews will be reviewed during a meeting September 19–23, 2022.

The **AOP** consisted of: Russell W. Brown, Ph. D.⁵ (chair), Gary Nelson, Ph. D.⁶, Lisa Kerr, Ph. D.⁷, Paul Rago, Ph. D.⁸ and Michael Wilberg, Ph. D.⁹.

Meeting Details:

These meetings were guided by the **NRCC** approved stock assessment guidance documents. Three background documents were provided to the Panel: (1) an updated prospectus for each stock; (2) an overview summary of all the salient data and model information for each stock; and (3) the **NRCC** Guidance memo on the Operational Assessments. Prior to the meeting, each assessment lead prepared a proposal for their Management Track Assessment. The proposal reflected the research track or most recent assessment results, the peer review panel Summary Report results and any initial investigations conducted for the management track assessment.

At the meeting, each assessment lead gave a presentation on the data to be used, model specifications (if applicable), evaluation of model performance, the process for updating the Biological Reference Points, the basis for catch projections, and an alternate assessment approach if their analytical assessment was rejected by the peer review panel.

Major Recommendations for Review of Individual Stocks:

In general, the **AOP** approved the plans presented, but recommended several points of emphasis to the recommended review levels as summarized below. **AOP** guidelines can be found in the stock assessment process document.

⁵Northeast Fisheries Science Center, Woods Hole, Massachusetts. (5/23, 5/24, 8/3)

⁶Atlantic States Marine Fisheries Commission, Massachusetts Division of Marine Fisheries. (5/23, 5/24, 8/3)

⁷Chair of the **NEFMC** Scientific and Statistical Committee, Gulf of Maine Research Institute. (5/23, 5/24, 8/3)

⁸Chair of the **MAFMC** Scientific and Statistical Committee, **NOAA** Fisheries (retired). (5/24, 8/3)

⁹Vice-chair of the **MAFMC** Scientific and Statistical Committee, University of Maryland. (5/23)

Table 2: Stocks reviewed at September 2022 Management Track Assessment Stock Assessments meeting

Stock	Lead	Review Level	Rationale and Comments
Ocean Pout	Charles Adams	Level 1 Direct Delivery	Little impact of swept area adjustments on survey indices, low biomass could result in variable survey indices. Concern about the reliability of the discards.
Atlantic Wolffish	Charles Adams	Level 1 Direct Delivery	Little impact of swept area adjustments on survey indices, low biomass could result in variable survey indices; AOP requested that the longline survey indices be presented with the assessment; some concern length frequency data.
Georges Bank Winter Flounder	Alex Hansell	Level 2 Expedited Review	VPA assessment update with little change to the previous assessment, major retrospective pattern in previous assessment.
Gulf of Maine Winter Flounder	Paul Nitschke	Level 2 Expedited Review	Little change proposed to the assessment methods, little impact of the swept area adjustments on the NEFSC survey indices; missing survey in 2020 is problematic and the panel would recommend investigating other averaging approaches.
Atlantic Halibut	Dan Hennen	Level 2 Expedited Review	There was a large increase in the 2020 Canadian catch attributable to increased targeting by their longline fishery. We encourage investigation of the U.S. longline and any Canadian indices of abundance. There is sufficient investigation planned in the assessment methods to warrant a Level 2 review.
Witch Flounder	Susan Wigley	Level 1 Direct Delivery	Witch Flounder currently uses an empirical approach to provide management advice. The NEFSC bottom trawl surveys will be updated to include swept area adjusted abundance and biomass surveys. 2020 survey values missing due to Covid will be treated as missing in the application of the empirical approach.
Monkfish North	Jon Deroba	Level 2 Expedited Review	Revise discard estimation method from simple ratio to combined ratio method to make this consistent with SBRM and other stocks. No longer manually remove some 'extreme' discard observations. Make the stat areas that define each management area consistent among landing and discard calculations. Reconsider discard mortality assumption of scallop dredge based on Weissman et al. 2021.
Monkfish South	Jon Deroba	Level 2 Expedited Review	Revise discard estimation method from simple ratio to combined ratio method to make this consistent with SBRM and other stocks. No longer manually remove some "extreme" discard observations. Make the stat areas that define each management area consistent among landing and discard calculations. Reconsider discard mortality assumption of scallop dredge based on Weissman et al. 2021.
Georges Bank Haddock	Liz Brooks	Level 2 Expedited Review	Research Track completed earlier in 2022. Swept area adjustments to survey indices are within the confidence bounds of the survey estimates. Some changes to the projection methodology vs. the VPA based projections done in agepro .
Gulf of Maine Haddock	Charles Perretti	Level 3 Enhanced Review	The analyst plans to follow up on recommendations from the Research Track peer review to include the Bottom Longline Survey as a survey index and to develop a quantitative model that accounts for cohort strength to replace the averaging of Weights-at-Age over recent years in the projection approach.
CCGM Yellowtail Flounder	Larry Alade	Level 1 Direct Delivery	Scheduled for a Research Track assessment in 2024. One of the last remaining VPA . Few changes will be implemented. Missing survey data were evaluated and concerned about the retrospective.
SNEMA Yellowtail Flounder	Chris Legault	Level 2 Expedited Review	Scheduled for a Research Track assessment in 2024. Catch and survey indices are likely to remain very low. The analyst plans to explore fleet selectivity and other aspects to tune the ASAP model and explore cold pool indices relative to current recruitment to determine appropriate years to use projections.
White Hake	Kathy Sosebee	Level 3 Enhanced Review	Update all fishery and survey data through 2021. For 2020 CAA , use pooled Age Length Key as done for half 2 2003 since no survey age data. Potentially add ASMFC shrimp survey (include new age data) and bottom longline surveys either using spring and fall bottom trawl ages or direct ages.
Atlantic Pollock	Brian Linton	Level 3 Enhanced Review	Use new area swept-based NEFSC spring and fall survey indices. Evaluate use of a historical recreational catch series (1970–1980). Use current ASAP base and sensitivity model configurations. Evaluate new assessment starting year of 1981. Evaluate use of a single fishery fleet (commercial + recreational). Update base and sensitivity model BRPs using 2010 SAW 50 SSB/SSB_{threshold} , F_{40%SPR} and SSB at F_{40%SPR} projection approach.
American Plaice	Larry Alade	Level 2 Expedited Review	A Research Track stock assessment for American Plaice was peer reviewed in July 2022 (less than 3 weeks before the AOP meeting). New sources of data included 2020 and 2021 landings estimated through the Catch Assessment and Monitoring System (CAMS) and discards will be derived using CAMS generated landings and discard ratios generated using recent observer data. Projections will be done internally within WHAM , which will result in internally consistent input data. The alternative approach will be an ASAP type model (without random effects) that can be implemented within the WHAM framework.

Individual Stock Discussion Summaries:

Ocean Pout (AOP Lead: Michael Wilberg)



Recommendation: Level 1 (Direct Delivery)

Ocean pout is assessed using the relative exploitation rate following the accepted assessment in the 2008 **GARM**, and its most recent status was overfished but overfishing not occurring. The assessment uses the catch divided by the 3-year moving average of the **NEFSC** Spring Trawl Survey. Catch is prohibited, so all catch is from discards. Projections are not done for this stock, and there is no alternate assessment approach. Recent years discards are estimated using the total **CAMS** catch and discard ratios from **NEFOP**. Recreational discards are not included as they are considered negligible. The survey indices had only very minor changes from the swept area adjustments. The stock appears to be at low biomass, which could result in variable survey indices.

The management track assessment will go through 2021. There are no proposed changes to analyses, but two data streams have changes in how they are calculated: the **NEFSC** Spring Trawl Survey Index and the discarded catch. These changes in processing the data streams are expected to have minor effects, but they could result in larger changes than anticipated. In particular, the lack of **NEFOP** sampling during part of 2020 has the potential to affect the estimate of discards. The direction and magnitude of that effect would depend on whether the period for which samples are not available is different from the rest of the period over which discards are calculated. Additionally, the 2020 **NEFSC** Spring Trawl Survey index is not available, and a two-year moving average will be used for the years impacted by that year. Therefore, the **AOP** recommends a Level 1 (Direct Delivery) review.

Atlantic Wolffish (AOP Lead: Lisa Kerr)




Recommendation: Level 1 (Direct Delivery)

Wolffish is currently assessed using a forward projection model known as **scale** (Statistical Catch At Length) which tunes to size and age data from trawl survey recruitment and adult indices, total catch, and catch size distributions along with overall growth information. The model was approved by the 2008 Northeast Data Poor Stocks Working Group (**NDPSWG**) and last updated in 2020. The model features one fishery fleet with one selectivity block and four survey indices (**NEFSC** spring recruitment, and **NEFSC** fall and spring and **MA DMF** adult indices). The most recent stock status was overfished but overfishing is not occurring. In 2010, there was a change to a no possession limit for wolffish, so catch is from discards since that time. The **NDPSWG** deemed projections unreliable for this stock assessment and they are not conducted.

The management track assessment will update data for this stock through 2021. There are no proposed changes to the model, but two data streams (i.e., **NEFSC** Trawl Survey and the discarded catch) have changes in how they are calculated and **Covid-19** disruptions resulted in missing surveys and reduced observer and port sampling of catch data in 2020. The **NEFSC** has adopted swept area biomass calculations of indices and the impact of the adjustment to the **NEFSC** trawl survey data was reported to be minimal for wolffish. The most recent years discards (2020 and 2021) will be estimated based on discard ratios calculated as usual based on observer data and the **CAMS** derived fleet landings. The missing 2020 survey

data will be treated as missing in the assessment (i.e., not imputed). The Panel expressed concern regarding the sufficiency of length frequency data in 2020 due to reduced observer coverage and how that could impact the assessment. The lead analyst noted that **scale** allows for missing length data and there are previous years with missing data. In addition, there is a general deficiency of data for this stock which would make it challenging to do much analytically about this issue. The **AOP** suggested that Bottom Longline Survey indices for wolffish be provided along with the assessment. The backup assessment approach is a 'Plan B' **Loess** smooth of **NEFSC** spring and fall adult indices. The uncertainty introduced by using ocean pout calibration and integration of newly published sex specific growth parameters were discussed but won't be addressed in this management track due to time limitations. This management track assessment will update data through 2021 and include a new survey index as additional information for consideration outside of the assessment. The **AOP** recommended a Level 1 (Direct Delivery) review for wolffish with the opportunity for an update from the analyst on any identified data or model issues at the August **AOP** meeting.

Georges Bank Winter Flounder (AOP Lead: Gary Nelson)




Recommendation: Level 2 (Expedited Review)

The current assessment method for Georges Bank Winter Flounder is a **VPA** model that includes age-specific **US** and Canadian landings and discards, and age-specific trawl indices (**NEFSC** fall, **NEFSC** spring and **CA DFO** spring surveys). The proposed work for the 2022 Management Track assessment includes updating all landings, discards and the survey data, and performing **OFL** and **ABC** projections at F_{MSY} for 2024–2026. The landings and discards will be updated via the **CAMS** system and the old **NEFSC** indices will be replaced with new **NEFSC** area-swept indices.

The **AOP** discussed the potential impact of the missing survey indices in 2020, data deficiencies with sampling and **CAMS** system estimates on the assessment. Comparisons between the new and old indices and between the 2019 **AA** and **CAMS** landings and discards were not provided, so members were uncomfortable concluding that the changes would have limited impacts on assessment results. In addition, the last **VPA** had a large retrospective bias and members expressed concern that low samples of data for characterizing catch-at-age could exacerbate the bias. Based on those concerns, the Panel elevated the assessment to a Level 2 (Expedited) review.

Gulf of Maine Winter Flounder (AOP Lead: Gary Nelson)



Recommendation: Level 2 (Expedited Review)

The current assessment method is an index-based approach that uses catch and estimates of 30+ **cm** biomass from three non-overlapping fall trawl surveys (**ME/NH**, **MA DMF**, **NEFSC**). The proposed work for the 2022 Management Track assessment is to update the fall surveys and catch through 2021. The plan proposes to replace the **NEFSC** standardized trawl index with the new area-swept adjusted index. In addition, catch data will be assembled using the new **CAMS** automated system.

The **AOP** discussed three primary issues with the proposed assessment update. The **AOP** was concerned about the impact that **Covid** restrictions in 2020 had on sampling activities (e.g., **MRIP** intercept

sampling, observer coverage, etc.) from which estimates of landings and discards are made, and they wondered what biases could occur in the estimations. The **AOP** discussed the impact of the missing survey indices in 2020 on the calculation of catch advice because it is usually computed by using the average of two recent fall surveys; therefore, members worried about potential instability in the updated estimate using only one year (2021). Also, the **AOP** conferred that, although differences between **NEFSC** new area-swept index and the old index appear minor, some unanticipated changes in the results could occur. Based on those discussions, the Panel agreed to elevate this assessment to a Level 2 (Expedited) review.

Atlantic Halibut (AOP Lead: Paul Rago)



Recommendation: Level 2 (Expedited Review)

Atlantic halibut catches limits are based on an index method that combines trends in several measures of relative abundance to adjust recent catches on a regular basis. The method is known as the First and Second Derivative (**FSD**) method because it adjusts catches using a linear combination of the slope and rate of change in slopes of abundance indices. The catch in year $t + 1$ is estimated as the product of the adjustment rate and the catch in year t . The abundance indices are the **NEFSC** fall bottom trawl survey and discard ratios for gillnet and trawl fleets in the Northeast. No biological reference points for Atlantic halibut in the **US** are available.

Application of the model in 2022 is complicated by a major change in the Canadian fishery in 2020. The total catch used in the original model includes catches in Canada from Stat Area 5. Shackell et al. (2021) reports the ongoing recovery of halibut but the recovery in Canadian waters may be occurring more quickly. Tagging analyses reported in Rago (2018) suggest regular movement of Atlantic halibut between **US** and Canada.

The sharp increase in landings in Canadian waters and declining indices in the **US** poses a dilemma for application of the current **FSD** model. Canada's increase in landings is driven by results of a **DFO** assessment that increased the quota. This assessment is likely to have indices that are trending upward in contrast to **US** indices which appear to be either level or slightly decreasing. Nonetheless, the slightly lower **FSD** multiplier, when multiplied by the increased total catch, results in a large increase in potential **US** catch. The appropriateness of this calculation was discussed but not resolvable during the **AOP** meeting.

The Panel suggested that an investigation of the basis for the increase in Canadian landings would be useful. Comparisons of **US** index trends with Canadian indices of abundance might also be useful. The assessment lead will also investigate the applicability of the Cooperative Longline survey in the Gulf of Maine in the **FSD** model. The assessment lead also proposes to modify and align some of the Stat Areas with survey areas but does not plan to redefine stock areas. In view of the potential changes in the model framework and addition of a new index, the Panel recommended a Level 2 (Expedited) review for Atlantic halibut.

Witch Flounder (AOP Lead: Russell Brown)



Recommendation: Level 1 (Direct Delivery)

Witch Flounder currently uses an empirical approach to provide management advice. It is a unit stock, so is less dependent on CAMS approaches to allocate catch to separate stock areas. The NEFSC bottom trawl surveys will be updated to include swept area adjusted abundance and biomass surveys. 2020 survey values missing due to Covid will be treated as missing in the application of the empirical approach. It was noted that the age structure of the population continues to be truncated and the analyst will include supplement data in the data portal that is not directly used in the empirical analysis. The panel concluded that a Level 1 (Direct Delivery) review was warranted.

Northern and Southern Monkfish (AOP Lead: Gary Nelson)



Recommendation: Level 2 (Expedited Review)

The current assessment method for the northern and southern Monkfish stocks is the index-based method known as ‘PlanBsmooth’ that uses fishery landings and discards, and NEFSC fall, spring and summer survey indices. The proposed work for the 2022 Management Track assessment includes updating all landings, discards and the survey data through 2021 (the spring survey will be updated through 2022). The landings will be updated via the CAMS system and a new method for estimating discards will be examined. Also, the old NEFSC indices will be replaced with new NEFSC area-swept indices and methods for dealing with the missing 2020 survey values will be explored. Additionally, the discard mortality assumption of Monkfish in scallop dredges will be re-examined, how extreme discard observations are handled will be changed, and adjustments to statistical areas that define the managements will be made consistent.

The main discussion of the AOP pertained to the proposed exploration of imputing missing survey values. One member wondered what the potential outcome would be and suggested that including an additional year further back in time might help with stability of resulting catch advice. The analyst responded that, based on earlier simulations examining biases in the ‘PlanBsmooth’ method, catch advice should be fairly robust with a missing year, but he will try the suggested method. The AOP panel agreed that a Level 2 (Expedited) review is appropriate for the proposed changes.


Georges Bank Haddock (AOP Lead: Russell Brown)



Recommendation: Level 2 (Expedited Review)

A Research Track Assessment for Haddock was completed earlier in 2022. The assessment will be updated through 2021 and utilize a WHAM state space model to develop estimates of recruitment, biomass, and fishery mortality. In the Research Track, the working group and analyst demonstrated extensive bridge building from VPA to ASAP, and from ASAP to WHAM (Research Track TOR #4). The panel was concerned that this is one of the first implementations of WHAM, uncertainty about the change in the scale of the catch advice, and about reduced sampling in recent years. On this basis, the panel recommended a Level 2 (Expedited) review for this stock.

Gulf of Maine Haddock (AOP Lead: Russell Brown)



Recommendation: Level 3 (Enhanced Review)

A Research Track Assessment for Haddock was completed earlier in 2022. The assessment will be updated through 2021 and utilize an ASAP model to develop estimates of recruitment, biomass, and fishery mortality. The analyst plans to follow up on recommendations from the Research Track peer review to include the Bottom Longline Survey as a survey index and to develop a quantitative model that accounts for cohort strength to replace the averaging of Weights-at-Age over recent years in the projection approach. Based on these proposed changes to the assessment methodology the AOP concurred with the analyst recommendation that the management track update should receive a Level 3 (Enhanced) review.

Cape Cod/Gulf of Maine Yellowtail Flounder (AOP Lead: Lisa Kerr)



Recommendation: Level 1 (Direct Delivery)

Cape Cod/Gulf of Maine (CCGM) yellowtail flounder is assessed using a VPA that was approved in 2008 at GARM III and was last updated in 2019. The model includes a single fishery fleet and fall and spring time series from three fishery independent surveys (NEFSC, MA DMF, and ME/NH trawl surveys). This assessment has retrospective issues and adjustments were made to the model results. The most recent stock status is not overfished and overfishing is not occurring.

Two data streams (i.e., NEFSC trawl survey and landings) have changes in how they are calculated and Covid-19 disruptions resulted in missing surveys and reduced observer and port sampling of catch data in 2020. The NEFSC has adopted swept area biomass calculations of indices and the impact of the adjustment to the NEFSC trawl survey data is minimal for CCGM yellowtail flounder. The transition from AA tables to CAMS is not anticipated to have a significant impact based on a 2019 data comparison. The missing survey data will be treated as missing in the assessment (i.e., not imputed). An initial analysis of the impact of missing survey data on the performance of the VPA suggests the impact will be minimal.

This management track assessment will update all fishery and survey data through 2021 and use the current VPA model configuration with no changes. Projections will be calculated and BRPs will be updated using the prescribed approach without changes. The analyst will perform a comparison of population size between the cooperative research twin trawl catchability study and the VPA model estimates. The alternative assessment is an empirical approach which applies catchability estimates from the twin trawl study to expand survey catch/tow to absolute biomass from Bigelow Spring and Fall survey estimates. There are no major changes to the assessment model or the types of data incorporated in the model. The analyses of impacts of changes in data streams and missing data suggest that these will have minimal impact. The AOP recommended a Level 1 (Direct Delivery) review for CCGM yellowtail flounder with the opportunity for an update on any identified data or model issues at the August AOP meeting. The last management track assessment cited concerns about the uncertainty and retrospective patterns in this stock assessment. This is one of the last remaining assessments that applies the VPA approach and it is scheduled for a Research Track Stock Assessment in 2024. Major changes will be addressed at that time with a likely change in model type.

Southern New England/Mid-Atlantic Yellowtail Flounder (AOP Lead: Paul Rago)



Recommendation: Level 2 (Expedited Review)

The current assessment model for **SNEMA** yellowtail is based on an **ASAP** model, accepted in 2012 at **SAW 54** and last updated in 2019. The model features an age dependent M , single fleet fishery, and three fishery independent surveys. Six selectivity blocks are used to model the stock from 1973 to 2018. The most recent selectivity block began in 2002. The stock is severely depleted, at about 10% of the $\frac{1}{2}B_{MSY}$ level, but overfishing is not occurring. Recruitment has been low and both reference points and projections are based on R estimates from 1990 onward.

The pandemic resulted in loss of both spring and fall bottom trawl surveys in 2020 and reductions in observer coverage. Funding issues reduced port sampling efforts. There are no recreational landings. Bottom trawl estimates will now use swept area per tow measurements to improve accuracy. Slight changes in overall means have been observed, but the variances of estimates tend to be large, overwhelming potential differences in scale. There does not seem to be any significant trend towards higher or lower values given adjustments for swept area. Landings estimates by stat area, previously based on the **AA** method, are expected to change only slightly as a result of the new **CAMS** approach. The new estimates of landings will not have any effects on estimates of discards but might be important in some instances in the future. Discard estimates are scaled by multiplying discard:kept ratios by total landings, which will change when **CAMS** rather than **AA** based estimates are used. The lead analyst has proposed to re-examine the selectivity blocks and other settings to improve model performance. The potential effects of swept area-adjusted survey indices will also be examined. Several recent publications in the literature have illustrated the utility of state-space models to estimate effects of environmental factors on stock dynamics. In particular, increases of the cold pool index (i.e., warmer) in the mid-Atlantic are associated with lower recruitment. These results cannot be directly included in the current model but they may be used to refine the range of years used to define ‘recent’ recruitment. Such a change, if justifiable, would alter both the biological reference points and abundance projections.

Given the potential effects of changes in selectivity blocks and the consideration of state space model results to inform the current assessment, the **AOP** recommended a Level 2 (Expedited) review for **SNEMA** yellowtail.

White Hake (AOP Lead: Lisa Kerr)



Recommendation: Level 3 (Enhanced Review)

White Hake is currently assessed using the **ASAP** model which was accepted in 2013 at **SAW 56** and was last updated in 2019. The model extends back to 1963 and includes one fishery fleet with two selectivity blocks and two trawl survey indices (**NEFSC** fall and spring). Catch at age information is not well characterized for this stock due to possible misidentification of species in the commercial and observer data, particularly in early years, low sampling of commercial landings in some years, and sparse discard length data. Pooled age length keys (**ALK**) have been used during periods with deficient age data. The current status is overfished and overfishing is not occurring. This assessment has retrospective issues and adjustments were made to the model results.

Two data streams (i.e., NEFSC Trawl Survey and landings) have changes in how they are calculated and Covid-19 disruptions resulted in missing surveys and reduced observer and port sampling of catch data in 2020. The NEFSC has adopted swept area biomass calculations of indices and the impact of the adjustment to the NEFSC trawl survey data will be reported for white hake. The impact of the transition from AA tables to CAMS for white hake will be documented based on a 2019 data comparison. The missing survey data will be treated as missing in the assessment (i.e., not imputed) and a pooled ALK will be used for 2020 CAA for commercial landings.

The management track assessment will update all fishery and survey data through 2021. In addition, two new indices will be considered in the management track, the ASMFC shrimp survey and the Bottom Longline Survey. The current ASAP model configuration will be used with the additional indices. The analyst will explore the model sensitivity to use of pooled ALK. The biological reference points will be updated using approach prescribed through SAW 56 and projections will be performed assuming catch in 2022 is equal to the PDT provided 2022 landings. The alternative assessment plan is Loess smoothing of both NEFSC surveys indices to infer future catch increase. Age information as an important source of uncertainty for this stock. Ageing was completed for white hake from the shrimp survey, however, there is still a need for observer ages and from the bottom longline survey.

This management track assessment will involve substantial changes, including the potential addition of a new survey index. The AOP agreed with the analyst's suggestion of a Level 3 (Enhanced) review for this stock.

Pollock (AOP Lead: Paul Rago)



Recommendation: Level 3 (Enhanced Review)

Pollock is currently assessed with an ASAP model that relies on dome shaped selectivity patterns for both the fishery and surveys. To ensure model convergence, the selectivity of oldest fish is fixed. The double dome model creates a "cryptic" biomass that cannot be estimated by survey data or captured by the commercial fishery.

Assessment scientists, managers, and even some harvesters have expressed concerns about the validity of the base (i.e., double-domed) model. An alternative model, which includes the same data but assumes a flat-top selectivity for survey indices, is used for comparison. Estimates of exploitable biomass from the base model compare favorably to the estimates from the sensitivity model (with a flat-top selectivity pattern for the survey).

The change from AA to CAMS appears to have little effect (0.1% increase) for 2019 comparison. Similarly, the use of a variable area per tow estimator appears to have little effect on the annual mean abundance estimates.

Commercial and recreational fisheries are modeled separately using data from 1970 to present. However, recreational catch before the start of the MRFSS program is assumed to be zero. To eliminate this inconsistency, a revised starting year of 1981 is proposed. The assessment lead also suggested pooling of recreational and commercial landings. This would eliminate the need to specify two selectivity patterns for these fleets. However, there may be some conflating of selectivity patterns because recreational catches

historically occurred inshore and on smaller fish (so called harbor pollock). The joint effects of changing the starting year, combining the recreational and commercial catches into a single fleet, and inherent instability of the base model are likely to require significant exploration of alternative model runs. After consideration of the number of changes, and their potential interactive effects, the Panel concurred with the lead scientist's recommendation for a Level 3 (Enhanced) review.

American Plaice (AOP Lead: Russell Brown)



Recommendation: Level 2 (Expedited Review)

The Research Track stock assessment for American Plaice was peer reviewed in July 2022 (less than 3 weeks before the AOP meeting). A state space WHAM model informed by data through 2019 was tabled by the Management Track and accepted by the CIE peer review panel. New sources of data included 2020 and 2021 landings estimated through the Catch Assessment and Monitoring System (CAMS) and discards will be derived using CAMS generated landings and discard ratios generated using recent observer data. Model diagnostics for the assessment through 2019 were well behaved and the retrospective pattern was relatively insignificant (no retrospective adjustments are anticipated). Projections will be done internally within WHAM, which will result in internally consistent input data. The alternative approach will be an ASAP type model (without random effects) that can be implemented within the WHAM framework. As a result of these proposed changes, the AOP concurred that this assessment should receive a Level 2 (Expedited) review.

AOP Meeting Conclusions:

The AOP met on May 23–24, 2022 to review the stock assessment plans for 14 stocks and on August 3, 2022 for one stock scheduled for the September 2022 Management Track cycle. The panel concluded that Level 1 reviews (Direct Delivery) were warranted for ocean pout, Atlantic wolffish, witch flounder, and Cape Cod/Gulf of Maine yellowtail flounder; Level 2 reviews (Expedited Review) for Georges Bank winter flounder, Gulf of Maine winter flounder, Atlantic halibut, northern and southern monkfish, Georges Bank haddock, Southern New England/Mid-Atlantic yellowtail flounder and American Plaice; and Level 3 reviews (Enhanced Review) for Gulf of Maine haddock, white hake, and pollock. The Level 2 and 3 reviews will occur during the September 2022 Management Track Peer Review scheduled for September 19–22, 2022. Changes in the required review level would be triggered by a Northeast Fisheries Science Center request to increase the review level for a given stock. The AOP could concur to increase the review level via email or request to reconvene the AOP panel to have further discussions with the stock assessment lead. Any need to reconvene the panel would be a publicly announced meeting and any subsequent changes to the review level would be publicized to assessment partners and stakeholders.

Appendix A.1. Meeting participants

Panel, May 2022:

Lisa Kerr – AOP (NEFMC SSC)
Gary Nelson – AOP (ASMFC)
Mike Wilberg and Paul Rago – AOP (MAFMC SSC)
Russ Brown – AOP Chair (NEFSC)
Michele Traver – NEFSC Assessment Process Lead

Attendees and Presenters, May 2022:

Alex Dunn – NEFSC
Alex Hansell – NEFSC
Andrew Jones – NEFSC
Angela Forristall – NEFMC
Benjamin Levy – NEFSC
Brian Linton – NEFSC
Cate O’Keefe – Fishery Applications Consultant
Charles Adams – NEFSC
Charles Perretti – NEFSC
Chris Kellogg – NEFMC
Chris Legault – NEFSC
Christopher Maguire – Nature Conservancy
Daniel Hennen – NEFSC
Gareth Lawson – Conservation Law Foundation
Jamie Cournane – NEFMC
Janice Plante – NEFMC
Jean-Jacques Maguire – NEFMC SSC
Jennifer Couture – NEFMC
John Pappalardo – NEFMC member, Cape Cod Commercial Fishermen’s Alliance
Jon Deroba – NEFSC
Julie Nieland – NEFSC
Katherine Sosebee – NEFSC
Kelley Whitmore – MA DMF
Kiersten Curti – NEFSC
Larry Alade – NEFSC
Libby Etrie – NEFMC member, Northeast Sector Service Network, Inc.
Liz Brooks – NEFSC
Liz Sullivan – GARFO
Maggie Raymond – Associated Fisheries of Maine
Melissa Sanderson – Cape Cod Commercial Fishermen’s Alliance
Paul Nitschke – NEFSC
Peter Melanson – Protech AIS
Rachel Feeney – NEFMC
Rebecca Peters – ME DMR

Rick Bellavance – NEFMC Vice Chair, RI Party and Charter Boat Association
Robin Frede – NEFMC
Spencer Talmage – GARFO
Steve Cadrin – SMAST
Susan Wigley – NEFSC
Tara Dolan – MADMF
Thomas Nies – NEFMC Executive Director
Tracey Bauer – ASMFC

Appendix B. Management Track Stock Assessment Terms of Reference

1. Estimate catch from all sources including landings and discards.
2. Evaluate indices used in the assessment (e.g., indices of relative or absolute abundance, recruitment, state surveys, age-length data, etc.).
3. Estimate annual fishing mortality, recruitment and stock biomass (both total and spawning stock) as possible (depending on the assessment method) for the time series using the approved assessment method and estimate their uncertainty. Include retrospective analyses if possible (both historical and within-model) to allow a comparison with previous assessment results and projections, and to examine model fit.
 - a. Include bridge runs to sequentially document each change from the previously accepted model to the updated model proposed for this peer review.
 - b. Prepare a backup assessment approach that would serve as an alternative for providing scientific advice to management if the analytical assessment were to not pass review.
4. Re-estimate or update the BRPs as defined by the management track level and recommend stock status. Also, provide qualitative descriptions of stock status based on simple indicators/metrics (e.g., age- and size-structure, temporal trends in population size or recruitment indices, etc.).
5. Conduct short-term stock projections when appropriate.
6. Respond to any review panel comments or SSC concerns from the most recent prior research or management track assessment.

Note: Major changes from the previous stock assessment require pre-approval by the Assessment Oversight Panel.

Appendix C. September 2022 Management Track Peer Review meeting attendees.

Panel, September 2022:

Richard Merrick – Chair
Matt Cieri – Panel
Cate O’Keefe – Panel
Yan Jiao – Panel
Russ Brown – NEFSC
Michele Traver – NEFSC

Attendees and Presenters, September 2022:

Alan d’Entremont– Scotia Harvest Inc., TMGC Canadian co-chair
Alex Dunn – NEFSC
Alex Hansell – NEFSC
Angela Forristall – NEFMC
Bill Devoe – ME DMR
Brian Linton – NEFSC
Carl Wilson – ME DMR
Charles Adams – NEFSC
Charles Perretti – NEFSC
Chris Kellogg – NEFMC
Chris Legault – NEFSC
Dan Hennen – NEFSC
Dave McElroy – NEFSC
Dave Richardson – NEFSC
Gareth Lawson – Conservation Law Foundation
Gary Nelson – MA DMF
Jackie O’Dell – Northeast Fisheries Coalition
Jamie Cournane – NEFMC
Jason Didden – MAFMC
Jennifer Couture – NEFMC
Jon Deroba – NEFSC
Jonathon Peros – NEFMC
Joseph Powers – NOAA (retired)
Julie Nieland – NEFSC
Kathy Sosebee – NEFSC
Kelly Whitmore – MA DMF
Kiersten Curti – NEFSC
Kris Vascotto – Atlantic Groundfish Council, Executive Director
Larry Alade – NEFSC
Libby Etrie – Northeast Sector Service Network, Inc.

Liz Brooks – NEFSC
Liz Sullivan – GARFO
Melanie Griffin – MA DMF
Mark Terceiro – NEFSC
Paul Nitschke – NEFSC
Rachel Feeney – NEFMC
Rebecca Peters – ME DMR
Rick Bellavance – RI Party and Charter Boat Association
Robin Frede – NEFMC
Spencer Talmage – GARFO
Steve Cadrin – SMAST
Susan Wigley – NEFSC
Tara Dolan – MA DMF
Tara Trinko Lake – NEFSC
Tom Miller – NEFSC
Tom Nies – NEFMC, Executive Director
Tracey Bauer – NCDMF
Xavier Mouy – NEFSC
Yanjun Wang – DFO



Aerial view of NMFS building and surrounds, Woods Hole Laboratory, MA; photo ©WHOI

Appendix D. Realized Agenda for September 2022 Management Track peer review

Time	Activity	Lead
Monday, September 19, 2022		
9:00–9:15 am	Welcome/Logistics/Conduct of Meeting	Michele Traver, Russ Brown, Richard Merrick, Chair
9:15–10:00 am	Input Data Changes Discussion/Questions	Russ Brown, Review Panel
10:00–11:00 am	GOM Winter flounder Discussion/Questions	Paul Nitschke, Review Panel
11:00–11:15 am	— Break —	
11:15–12:15 pm	George Bank winter flounder Discussion/Questions	Alex Hansell, Review Panel
12:15–12:30 pm	Discussion/Summary	Review Panel
12:30–12:45 pm	Public Comment	Public
12:45–1:45 pm	— Lunch —	
1:45–2:45 pm	Atlantic halibut Discussion/Questions	Dan Hennen, Review Panel
2:45–3:45 pm	Georges Bank haddock Discussion/Questions	Liz Brooks, Review Panel
3:45–4:00 pm	— Break —	
4:00–4:15 pm	Discussion/Summary	Review Panel
4:15–4:30 pm	Public Comment	Public
4:30 pm	— Adjourn —	
Tuesday, September 20, 2022		
9:00–9:05 am	Welcome/Logistics	Michele Traver, Richard Merrick, Chair
9:05–10:30 am	White hake	Kathy Sosebee
10:30–10:45 am	— Break —	
10:45–12:00 am	White hake cont. Discussion/Questions	Kathy Sosebee, Review Panel
12:00–12:15 pm	Discussion/Summary	Review Panel
12:15–12:30 pm	Public Comment	Public
12:30–1:30 pm	— Lunch —	
1:30–3:30 pm	Monkfish (North and South) Discussion/Questions	Jon Deroba, Review Panel
3:30–3:45 pm	— Break —	
3:45–4:45 pm	Southern New England/Mid-Atlantic yellowtail flounder Discussion/Questions	Chris Legault
4:45–5:00 pm	Discussion/Summary	Review Panel
5:00–5:15 pm	Public Comment	Public
5:15 pm	— Adjourn —	
Wednesday, September 21		
9:00–9:05 am	Welcome/Logistics	Michele Traver, Richard Merrick, Chair
9:05–10:30 am	Gulf of Maine haddock	Charles Perretti
10:30–10:45 am	— Break —	
10:45–12:00 pm	Gulf of Maine haddock cont. Discussion/Questions	Charles Perretti, Review Panel
12:00–12:15 pm	Discussion/Summary	Review Panel
12:15–12:30 pm	Public Comment	Public
12:30–1:30 pm	— Lunch —	
1:30–3:30 pm	Pollock	Brian Linton
3:30–3:45 pm	— Break —	
3:45–4:45 pm	Pollock cont. Discussion/Questions	Brian Linton
4:45–5:00 pm	Discussion/Summary	Review Panel
5:00–5:15 pm	Public Comment	Public
5:15 pm	— Adjourn —	
Thursday, September 22		
9:30–11:00 am	American plaice Discussion/Questions	Larry Alade
11:00–11:15 am	Discussion/Summary	Review Panel
11:15–11:30 am	Public Comment	Public
11:30–12:00 am	Key Points/Follow ups	Review Panel
12:00–1:00 pm	— Lunch —	
1:00–5:00 pm	Report Writing	Review Panel

2. ATLANTIC WOLFFISH

Charles Adams

This assessment of the Atlantic wolffish (*Anarhichas lupus*) stock is a Level 1 management track assessment of the existing benchmark assessment (NDPSWG 2009). Based on the previous 2020 management track assessment (NEFSC 2022) the stock was overfished, but overfishing was not occurring. This assessment updates commercial fishery catch data, research survey indices of abundance, the analytical scale assessment model and reference points through 2021.

State of Stock: Based on this updated assessment, the Atlantic wolffish (*Anarhichas lupus*) stock is overfished and overfishing is not occurring (Figures 1–2). Retrospective adjustments were not made to the model results. Spawning stock biomass (*SSB*) in 2021 was estimated to be 690 (mt) which is 46% of the biomass target ($SSB_{MSY\ proxy} = 1,509$; Figure 1). The 2021 fully selected fishing mortality was estimated to be 0.004 which is 2% of the overfishing threshold proxy ($F_{MSY\ proxy} = 0.192$; Figure 2).

Table 3: Catch and status table for Atlantic wolffish. All weights are in (mt), recruitment is in (millions) and F_{Full} is the fully selected fishing mortality. Model results are from the current updated scale assessment.

	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
	<i>Data</i>									
Commercial landings	0	0	0	0	0	0	0	0	0	0
Commercial discards	3	2	1	1	1	2	3	3	2	2
Recreational landings	0	0	0	0	0	0	0	0	0	0
Catch for Assessment	3	2	1	1	1	2	3	3	2	2
	<i>Model Results</i>									
Spawning Stock Biomass	368	424	476	522	567	607	638	660	674	690
F_{Full}	0.01	0.005	0.003	0.003	0.002	0.005	0.005	0.005	0.004	0.004
Recruits (age-1)	50	45	39	56	96	138	273	274	274	274

Table 4: Comparison of reference points estimated in the 2020 management track and from the current assessment update. An $F_{40\%SPR}$ proxy was used for the overfishing threshold and was based on yield per recruit calculations within the scale model.

	2020	2022
$F_{MSY\ proxy}$	0.200	0.192
SSB_{MSY} (mt)	1,543	1,509
<i>MSY</i> (mt)	218	211
Median recruits (age-1) (millions)	238	232
<i>Overfishing</i>	No	No
<i>Overfished</i>	Yes	Yes

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

The primary sources of uncertainty are the use of the ocean pout calibration coefficient (Atlantic wolffish coefficients are unknown), and the change to a no possession limit in May 2010. The ocean pout calibration coefficient (4.575) is one of the largest for any species (Miller et al. 2010), and results in lower biomass estimates. The change to a no possession limit places greater importance on discard mortality. Additionally, it is unclear whether the lack of a recruitment index since 2005 is due to an actual decrease in recruitment, or a change in catchability resulting from the increase in liner mesh size associated with the switch to the Bigelow. Other sources of uncertainty were identified in previous Atlantic wolffish assessments (NDPSWG 2009, NEFSC 2012): the surveys may have reached the limit of wolffish detectability due to the decline in abundance; and the lack of commercial length information results in model estimation difficulties for fishery selectivity.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted SSB or F_{Full} lies outside of the approximate joint confidence region for SSB and F_{Full}).

This assessment has retrospective patterns with Mohn's $\rho = 0.18$ for SSB and -0.10 for F . However, confidence intervals are not available because MCMC is not fully developed for the scale model.

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?

Due to the uncertainties in the assessment, the Northeast Data Poor Stocks Working Group (NDPSWG 2009) concluded that stock projections would be unreliable and should not be conducted. Catch advice is derived as $OFL = F_{MSY} \times B_{Terminal}$ using the terminal year exploitable biomass.

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the effect these changes had on the assessment and stock status.

The time series of Bigelow indices was recalculated using station-specific swept areas. Supplemental Figure 26 (see SASINF) was presented to the Assessment Oversight Panel (AOP) on May 23, 2022; the AOP agreed that the differences were minor.

The data source for commercial landings changed to the Catch Accounting and Monitoring System (CAMS) beginning in 2020. However, given the no possession limit, the AOP agreed that this is not an issue.

- If the stock status has changed a lot since the previous assessment, explain why this occurred.

Stock status has not changed since the previous assessment.

- Provide qualitative statements describing the condition of the stock that relate to stock status.

Catch has been limited almost exclusively to discards since the implementation of the no possession rule in May 2010. No age-1 recruits have been caught in the NEFSC spring survey since 2005.

- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.

Several research needs were identified by the Peer Review Panel in the 2015 assessment (NEFSC 2015): potential use of a likelihood profile to apply the criterion for a retrospective adjustment; further studies on growth parameters; a tagging study to provide information on stock structure and movement; and a study of post-capture nest site fidelity.

- Are there other important issues?

All 2020 surveys were treated as missing in the [scale](#) model. However, it should be noted that the [scale](#) model treats survey indices with zero catch as missing as well. For example, the [NEFSC](#) spring adult index had zero catch in 2004–2006, 2008 and 2011; thus, these years are treated as missing by the [scale](#) model.

Recruitment at the end of the time series increases toward the initial recruitment estimate (Table 3; Figure 3) because there is no information in the model to inform these estimates. There is no indication in the data that recruitment has increased recently.

Approximate 90% [log-normal](#) confidence intervals are not shown in Figures 1–3 because [MCMC](#) is not fully developed for the [scale](#) model.

Discards estimates assume an 8% mortality rate based on Grant and Hiscock (2014). This results in very low removals under the no possession rule. Future model updates should see a population response from these low removals. However, if no change is observed in the data inputs (e.g., increased survey indices) then the diagnostics may worsen.

Bottom long-line survey indices, which are not currently used in the [scale](#) model, are shown in supplemental Figure 27 (see [SASINF](#)) for informational purposes.



Wolffish at floor of aquarium tank. Photo credit: Woods Hole Aquarium

2.1. Reviewer Comments: Atlantic wolffish

Atlantic wolffish was not peer reviewed in fall of 2022.

References:

Grant S.M., Hiscock W. 2014. Post-capture survival of Atlantic wolffish (*Anarhichas lupus*) captured by bottom otter trawl: Can live release programs contribute to the recovery of species at risk? *Fish Res* 151:169–176. [11.003](#)

Miller T.J., Das C., Politis P.J., Miller A.S., Lucey S.M., Legault C.M., Brown R.W., Rago P.J. 2010. Estimation of Albatross IV to Henry B. Bigelow calibration factors. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 10-05; 233p.

Northeast Data Poor Stocks Working Group ([NDPSWG](#)). 2009. The Northeast Data Poor Stocks Working Group Report, December 8–12, 2008 Meeting. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 09-02; 496p.

Northeast Fisheries Science Center ([NEFSC](#)). 2012. Assessment or data updates of 13 Northeast groundfish stocks through 2010. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 12-06; 789p. [CRD12-06](#)

Northeast Fisheries Science Center ([NEFSC](#)). 2015. Operational assessment of 20 Northeast groundfish stocks, Updated Through 2014. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 15-24; 251p. [CRD15-24](#)

Northeast Fisheries Science Center ([NEFSC](#)). 2022. Fall management track assessments 2020. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 22-08; 168p. [CRD22-08](#)



Anarhichas lupus, Atlantic wolffish.




Figure 1: Trends in spawning stock biomass of Atlantic wolffish between 1968 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $SSB_{Threshold}$ ($\frac{1}{2}SSB_{MSY proxy}$; horizontal dashed line) as well as SSB_{Target} ($SSB_{MSY proxy}$; horizontal dotted line) based on the 2022 assessment. Biomass was not adjusted for a retrospective pattern.




Figure 2: Trends in the fully selected fishing mortality (F_{Full}) of Atlantic wolffish between 1968 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $F_{\text{Threshold}}$ ($F_{\text{MSY proxy}} = 0.192$; horizontal dashed line) based on the 2022 assessment. F_{Full} was not adjusted for a retrospective pattern




Figure 3: Trends in age-1 recruits (millions) of Atlantic wolffish between 1968 and 2021 from the current (solid line) and previous (dashed line) assessment.




Figure 4: Total catch of Atlantic wolffish between 1968 and 2021 by fleet (commercial and recreational) and disposition (landings and discards). Note that a no possession limit was put in place in May 2010.




Figure 5: Indices of biomass for Atlantic wolffish between 1968 and 2021 for the Northeast Fisheries Science Center (NEFSC) spring and fall bottom trawl surveys, and the Massachusetts Division of Marine Fisheries (MADMF) spring bottom trawl survey. The approximate 90% log-normal confidence intervals are shown.

3. GULF OF MAINE WINTER FLOUNDER

Paul Nitschke

This assessment of the Gulf of Maine winter flounder (*Pseudopleuronectes americanus*) stock is a Management Track assessment of the existing 2020 area-swept Management Track assessment (NEFSC 2022). Based on the previous assessment the biomass status is unknown but overfishing was not occurring. This assessment updates commercial and recreational fishery catch data, research survey indices of abundance, and the area-swept estimates of 30+ cm biomass based on the fall NEFSC, MA DMF, and ME/NH surveys.

State of Stock: Based on this updated assessment, the Gulf of Maine winter flounder (*Pseudopleuronectes americanus*) stock biomass status is unknown and overfishing is not occurring (Figures 6–7). Retrospective adjustments were not made to the model results. Biomass (30+ cm mt) in 2021 was estimated to be 5,093 mt (Figure 6). The 2021 30+ cm exploitation rate was estimated to be 0.033 which is 14% of the overfishing exploitation threshold proxy ($E_{MSY proxy} = 0.23$; Figure 7).

Table 5: Catch and status table for Gulf of Maine winter flounder. All weights are in (mt) and E_{Full} is the exploitation rate on 30+ cm fish. Biomass is estimated from survey area-swept for non-overlapping strata from three different fall surveys (ME/NH, MA DMF, NEFSC) using an updated q estimate of 0.81 based on the wing spread from the sweep study (Miller et al., 2017).

	2016	2017	2018	2019	2020	2021
<i>Data</i>						
Recreational discards	11	5	2	2	1	1
Recreational landings	41	161	80	42	51	43
Commercial discards	3	3	3	4	2	6
Commercial landings	185	210	158	102	81	118
Catch for Assessment	240	378	243	150	134	168
<i>Model Results</i>						
30+ cm Biomass	3,037	3,039	2,610	2,620	NA	5,093
E_{Full}	0.079	0.124	0.093	0.057		0.033

Table 6: Comparison of reference points estimated in an earlier assessment and from the current assessment update. An $E_{40\%}$ exploitation rate proxy was used for the overfishing threshold and was based on a length based yield per recruit model from the 2011 SARC 52 benchmark assessment.

	2020	2022
$E_{MSY proxy}$	0.23	0.23
B_{MSY}	Unknown	Unknown
MSY (mt)	Unknown	Unknown
Overfishing	No	No
Overfished	Unknown	Unknown

Projections: Projections are not possible with area-swept based assessments. Catch advice was based on 75% of $E_{40\%}$ ($75\% E_{MSY\ proxy}$) using the terminal year fall area-swept estimate assuming $q = 0.81$ on the wing spread which was updated using the average efficiency from 2009–2021 from the sweep experiment (Miller et al., 2017). Updated 2021 fall 30+cm area-swept biomass (5,093 mt) implies an OFL of 1,171 mt based on the $E_{MSY\ proxy}$ and a catch of 879 mt for 75% of the $E_{MSY\ proxy}$. Catch advice (OFLs and ABCs) from the 2020 Management Track assessment was based on the average of the last two years of the fall surveys to make better use of the available new information and to help stabilize the catch advice. Alternatively, since the 2020 surveys are not available due to Covid, using the average of updated 2021 and 2022 spring and 2021 fall 30+cm area-swept biomass (4,660 mt) implies an OFL of 1,072 mt based on the $E_{MSY\ proxy}$ and a catch of 804 mt for 75% of the $E_{MSY\ proxy}$.

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

The largest source of uncertainty with the direct estimates of stock biomass from survey area-swept estimates originates from the survey gear catchability (q). Biomass and exploitation rate estimates are sensitive to the survey q assumption. However this 2022 update does incorporate the use of a re-estimated q through an average estimate of efficiency from 2009–2021 fall and 2009–2022 spring ($q = 0.81$ fall and $q = 0.70$ spring) from the sweep study for the NEFSC survey. This updated q assumption (0.81) results in a lower estimate of 30+cm biomass (5,093 mt) relative to the 2020 estimate $q = 0.71$ assumption (5,783 mt) from the updated fall surveys. Another major source of uncertainty with this method is that biomass based reference points cannot be determined and overfished status is unknown.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted SSB or F_{Full} lies outside of the approximate joint confidence region for SSB and F_{Full} .)

The model used to determine status of this stock does not allow estimation of a retrospective pattern. An analytical stock assessment model does not exist for Gulf of Maine winter flounder. An analytical model was no longer used for stock status determination at SARC 52 (2011) due to concerns with a strong retrospective pattern. Models have difficulty with the apparent lack of a relationship between a large decrease in the catch with little change in the indices and age and/or size structure over time.

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?

Population projections for Gulf of Maine winter flounder do not exist for area-swept assessments and stock biomass status is unknown. This stock was never declared as overfished. Catch advice from area-swept estimates tend to vary with inter-annual variability in the surveys. Consideration was given to using multiple surveys (fall 2021 and spring 2021–2022) to stabilize the biomass estimates and catch advice since 2020 surveys are not available due to Covid.

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the effect these changes had on the assessment and stock status.
The assumption on q changed from 0.71 to 0.81 for the fall and from 0.62 to 0.70 for the spring using information from the updated average qs from the NEFSC survey (Miller et al., 2017) and incorporation of new survey data were made to this Gulf of Maine winter flounder Management Track assessment. The 2020 and 2021 commercial catch estimates are based on CAMS in this assessment. However, changes in total removals will not directly affect the estimated biomass or catch advice and total removals still remain far below the overfishing definition. In addition there were some minor changes to the survey indices due to tow based area-swept adjustments.
- If the stock status has changed a lot since the previous assessment, explain why this occurred.
The overfishing status of Gulf of Maine winter flounder has not changed.
- Provide qualitative statements describing the condition of the stock that relate to stock status.
The Gulf of Maine winter flounder has relatively flat survey indices with little change in the size structure over time. There have been large declines in the commercial and recreational removals since the 1980s. This large decline over the time series does not appear to have resulted in a response in the stock's size structure within the catch and surveys nor has it resulted in a change in the survey indices of abundance. However, there have been increases in the fall 2021 and the spring 2021 and 2022 area swept biomass estimates. If increasing biomass trends continue then perhaps this is the beginning of a response to time series lows in exploitation rates.
- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.
Direct area-swept assessments could be improved with additional studies on state survey gear efficiency. Quantifying the degree of herding between the doors and escapement under the footrope and/or above the headrope for state surveys is needed to improve the area-swept biomass estimates. Studies quantifying winter flounder abundance and distribution among habitat types and within estuaries could improve the biomass estimate.
- Are there other important issues?
The general lack of a response in survey indices and age/size structure are the primary sources of concern with catches remaining far below the overfishing level. Recent increases in the biomass could perhaps be the being of a response to removals being at record lows over the last three years (2019–2021). If recent increases in biomass is a response to the low catches then continuation of keeping catches near recent levels should result in further increases in biomass.

3.1. Reviewer Comments: Gulf of Maine winter flounder

The 2022 assessment of Gulf of Maine winter flounder (*Pseudopleuronectes americanus*) updates the 2020 area-swept Management Track assessment (NEFSC 2022)¹⁰. The analytic method was rejected in 2008 with GARM (2008) and again at SARC 52 (2011). Area swept assessments have been used since then. Updates were made of commercial and recreational fishery catch data, research survey indices of abundance, and the area-swept estimates of 30+cm biomass based on the fall NEFSC, Massachusetts Department of Marine Fisheries (MADMF), and Maine/New Hampshire (ME/NH) surveys.

Retrospective adjustments were not made to the model results. A small change in q of the NEFSC surveys resulted from a recent paired tow efficiency study (Miller et al. in 2020)¹¹ which reduced the biomass estimate for much of the time series. Biomass (30+cm) in 2021 was estimated to be 5,093 mt.

The 2021 30+cm exploitation rate was estimated to be 0.033 which is 14% of the overfishing exploitation threshold proxy ($E_{MSY\ proxy} = 0.23$).

Based on this updated assessment, the Gulf of Maine winter flounder stock's overfished status is unknown but overfishing is not occurring.

Projections are not possible with area-swept based assessments. However, catch advice can still be provided, and as suggested by the AOP, the assessment scientist considered two approaches:

- One approach provided catch advice using a method like that used with the 2020 Operational Assessment, which averaged the last two years of the fall surveys to make better use of the available new information and to help stabilize the catch advice. Because the 2020 fall survey was not conducted, the updated 2021 and 2022 spring and 2021 fall 30+cm area-swept biomass were averaged (4,660mt) which implies an OFL of 1,072mt based on the $E_{MSY\ proxy}$ and a catch of 804mt for 75% of the $E_{MSY\ proxy}$.
- A second approach provided was to base the catch advice on 75% of $E_{40\%}$ (75% $E_{MSY\ proxy}$) using the terminal year fall survey area-swept estimate, assuming $q = 0.81$ on the wing spread. The latter was updated using the average efficiency from 2009–2021 from the sweep experiment (Miller et al. 2020). Updated 2021 fall 30+cm area-swept biomass (5,093 mt) implies an OFL of 1,171 mt based on the $E_{MSY\ proxy}$ and a catch of 879 mt for 75% of the $E_{MSY\ proxy}$.

The Peer Review Panel (Panel) recommended that while the choice of approaches to providing catch advice does not have a major impact, it was the Panel's consensus that averaging the 2 spring (2021 and 2022) and 1 fall (2021) surveys was the better choice for this assessment. Averaging reduces the noise resulting from 'year' effects (i.e., the balance between day and night tows, plus length effects in the

¹⁰NEFSC. 2022. Fall Management Track Assessments 2020., US Dept Commer, Northeast Fish Sci Cent Ref Doc. 22-08; 168p. [CRD22-08](#)

¹¹Miller, T.J., D. Richardson, P. Politis, J. Blaylock, J. Manderson, and C. Roebuck. 2020. Relative efficiency of a chain sweep and the rockhopper sweep used for the NEFSC bottom trawl survey and biomass estimates for winter and windowpane flounder and red hake stocks. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 20-XX; 31p. [CRD20-XX](#)

daytime tows). The protocol for the next (e.g., 2024) assessment will need to revisit this decision once multiple sequential fall surveys are available.

The Panel is concerned more about the uncertainty surrounding the rapid increase in catch advice given the stock's depressed condition despite low fishing pressure. These historically low exploitation rates could be leading to the increased projected abundance seen in the most recent surveys, and an increase in effort could cap the stock's nascent recovery.

Research suggestions:

- The Center should also consider statistical approaches that overcome the imbalance between day and night tows in a stratum.
- Consider applying year specific qs rather than averaging the full time series.

The Panel concluded that the 2022 assessment update for Gulf of Maine winter flounder fulfilled the recommendations of the AOP, is technically sufficient to partially evaluate stock status and provide scientific advice and meets the Terms of Reference for the stock's assessment. The assessment represents Best Scientific Information Available (BSIA) for this stock for management purposes.

References:

Northeast Fisheries Science Center. 2022. Fall Management Track Assessments 2020, US Dept Commer, Northeast Fish Sci Cent Ref Doc. 22-08; 168p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026. [CRD22-08](#)

Northeast Fisheries Science Center. 2011. 52nd Northeast Regional Stock Assessment Workshop (SAW 52) Assessment Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 11-17; 962p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026. [CRD11-17](#).



Pseudopleuronectes americanus, Winter Flounder.




Figure 6: Trends in 30+cm area-swept biomass of Gulf of Maine winter flounder between 2009 and 2021 from the current assessment based on the fall (ME/NH, MA DMF, NEFSC) surveys.




Figure 7: Trends in the exploitation rates (E_{Full}) of Gulf of Maine winter flounder between 2009 and 2021 from the current assessment based on the fall (ME/NH, MA DMF, NEFSC) surveys and the corresponding $F_{Threshold}$ ($E_{MSY proxy} = 0.23$; horizontal dashed line).




Figure 8: Total catch of Gulf of Maine winter flounder between 2009 and 2021 by fleet (commercial and recreational) and disposition (landings and discards). A 15% mortality rate is assumed on recreational discards and a 50% mortality rate on commercial discards.




Figure 9: Indices of biomass for the Gulf of Maine winter flounder between 1978 and 2022 for the Northeast Fisheries Science Center (NEFSC), Massachusetts Division of Marine Fisheries (MADMF), and the Maine New Hampshire (ME/NH) spring and fall bottom trawl (strata 1–3) surveys. NEFSC indices are calculated with gear and vessel conversion factors where appropriate. The approximate 90% log-normal confidence intervals are shown.

4. GEORGES BANK WINTER FLOUNDER

Alex Hansell

This assessment of the Georges Bank Winter Flounder (*Pseudopleuronectes americanus*) stock is a Management Track update of the existing 2020 operational VPA assessment which included data for 1982–2020 (NEFSC 2020). Based on the previous assessment the stock was overfished and overfishing was not occurring. This assessment updates commercial fishery catch data, research survey biomass indices, and the analytical VPA assessment model and reference points through 2021. Additionally, stock projections have been updated through 2025.

State of Stock: Based on this updated assessment, the Georges Bank Winter Flounder (*Pseudopleuronectes americanus*) stock is not overfished and overfishing is not occurring (Figures 10–11). Retrospective adjustments were made to the model results. Spawning stock biomass (*SSB*) in 2021 was estimated to be 7159 mt. The 2021 fully selected fishing mortality (*F*) was estimated to be 0.0485. However, the 2021 point estimate of *SSB* and *F*, when adjusted for retrospective error (59% for *SSB* and –36% for *F*), are outside the 90% confidence intervals of the unadjusted 2021 point estimates. Therefore, the values used in the stock status determination were the retrospective-adjusted values of $F_{2021} = 0.076$ which is 16% of the overfishing threshold ($F_{MSY\ proxy} = 0.452$; Figure 11), and $SSB_{2021} = 4,503\text{ mt}$ which is 60% of the biomass target ($SSB_{MSY\ proxy} = 7,503$ with a threshold of 50% of $SSB_{MSY\ proxy}$; Figure 10).

Table 7: Catch input data and VPA model results for Georges Bank Winter Flounder. All weights are in (mt), recruitment is in (000s) and F_{Full} is the fishing mortality on fully selected ages (ages 4–6). Catch and model results are only for the most recent years (2012–2021) of the current updated VPA assessment.

	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
	<i>Data</i>									
US landings	1,911	1,675	1,114	866	462	364	416	280	292	249
CA landings	83	12	12	13	4	6	9	11	7	6
US discards	126	46	46	19	5	14	41	20	49	6
CA scall dr discards	79	28	47	42	21	16	22	18	49	22
Catch for Assessment	2,199	1,761	1,219	940	492	400	488	329	397	283
	<i>Model Results</i>									
Spawning Stock Biomass	4,289	3,477	3,229	3,162	2,736	2,385	2,515	3,326	4,079	7,159
F_{Full}	0.5404	0.5693	0.4899	0.2499	0.2015	0.1526	0.231	0.1412	0.0917	0.0485
Recruits (age-1)	3,366	2,409	2,863	1,322	3,034	3,495	5,617	4,940	10,740	8,920

Projections: Short-term projections of biomass were derived by sampling from a cumulative distribution function of recruitment estimates (1982–2020 Y_C) from the final run of the adapt VPA model. The annual fishery selectivity, maturity ogive (a 3-year moving window), and mean weights-at-age used in the projection are the most recent five-year averages (2017–2021). An *SSB* retrospective adjustment factor of 0.629 was applied in the projections.

Table 8: Comparison of reference points estimated in the 2020 assessment and the current assessment update and stock status during 2021 and 2021, respectively. A proxy for F_{MSY} ($F_{40\%SPR}$) was used for the overfishing threshold and was based on long-term stochastic projections of the stock based on the 2017–2021 means for selectivity-, maturity- and mean weights-at-age, and a CDF of estimated recruitments (using the entire time series). $SSB_{MSY proxy}$ was used as the biomass target and was based on long-term stochastic projections of the stock fished at $F_{40\%SPR}$.

	2020	2022
$F_{MSY proxy}$	0.358	0.452
$SSB_{MSY proxy}$ (mt)	7,267 (4,143–11,113)	7,503 (4,790–10,705)
MSY (mt)	2,573 (1,520–3,835)	2,757 (1,811–3,918)
Median recruits (age-1) (000s)	8,470	8,759
Overfishing	No	No
Overfished	Yes	No

Table 9: Short-term projections of catch (mt) and spawning stock biomass (mt) for Georges Bank Winter Flounder based on a harvest scenario of fishing at $F_{MSY proxy}$ between 2023 and 2025. Catch in 2022 was estimated to be 278 (mt) by the Groundfish Plan Development Team.

Year	Catch (mt)	SSB (mt)	F_{Full}
2022	278	5,755 (4,427–7,445)	0.058
Year	Catch (mt)	SSB (mt)	F_{Full}
2023	2,360	6,322 (4,651–8,890)	0.452
2024	1,963	4,738 (3,053–5,247)	0.452
2025	1,819	5,236 (3,467–7,957)	0.452

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

The largest source of uncertainty is probably the estimate of natural mortality, which is based on longevity (max. age = 20). Natural mortality is not well studied in Georges Bank Winter Flounder and is assumed to be constant over time. Natural mortality affects the scale of the biomass and fishing mortality estimates.

VPA assumes catch is known without error, which in the case of Georges Bank Winter Flounder is certainly not true. Discards from the Canadian bottom trawl fleet were not provided by DFO and the precision of the Canadian scallop dredge discard estimates are uncertain. In addition, there are no length or age composition data for the Canadian landings or discards of GB winter flounder. The lack of age data for the Canadian spring survey catches requires the use of the US spring survey A/L keys for several disparate data streams, including the Canadian scallop discards, US otter trawl and scallop discards, despite selectivity differences. Various other gaps in catch data at age or length have been filled using decisions based on expert opinion and are difficult, if not impossible, to reproduce. Different decisions produce different model inputs and result in different

outcomes. The direction and magnitude of the bias associated with filling gaps using expert opinion is unknown, but likely common in VPA assessments.

Another potentially important uncertainty is the lack of 2020 NEFSC fall and spring surveys. For 2022, DFO survey estimates were not available due to the use of a new survey vessel and an absence of a calibration factor.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted SSB or F_{Full} lies outside of the 90% confidence intervals for SSB and F_{Full} .)

The 7-year Mohn's ρ , relative to SSB , was 0.57 in the 2020 assessment and was 0.59 in 2021. The 7-year Mohn's ρ , relative to F , was -0.34 in the 2020 assessment and was -0.36 in 2021. There was a major retrospective pattern for this assessment because the ρ -adjusted estimates of 2021 SSB ($SSB_{\rho} = 4,503$) and 2021 F ($F_{\rho} = 0.076$) were outside the 90% confidence limits for SSB (6,871–11,642) and F (0.03–0.049). A retrospective adjustment was made for both the determination of stock status and for projections. The retrospective adjustment changed the 2021 SSB from 7,159 to 4,503 and the 2021 F_{Full} from 0.0485 to 0.076.

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?

Population projections for Georges Bank Winter Flounder are uncertain because confidence bounds for projected biomass estimates from the previous assessment did not capture the terminal estimate of biomass from this one. This stock was required to be rebuilt by 2017, but this did not occur. The stock is in a revised rebuilding plan, based on fishing at 70% of $F_{MSY proxy}$, with rebuilding by 2029.

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the affect these changes had on the assessment and stock status.

Changes made to the Georges Bank Winter Flounder assessment included updating the most recent 5-year averages (2017–2021) of fishery selectivity-, proportion mature-, stock weights-, catch weights-, and spawning stock weights-at-age.

US spring and fall indices were revised from 2009 to 2022 to account for tow-specific area swept; revised indices were similar to previous relative abundance estimates. The new US commercial fishery data processing system (Catch Accounting and Monitoring System [CAMS]) was used to produce US landings estimates for 2020 and 2021. In 2021, age samples were not available for US landings so the A/L key from the 2020 landings were used to produce estimates for 2021. The Covid epidemic caused the cancelation of the 2020 NEFSC spring and fall surveys. DFO survey results were not available for 2022 because of a change in vessel. The missing NEFSC surveys appear to have had a minor effect; however, the absence of the DFO survey most likely increased SSB estimates. The DFO survey results will be available once a calibration study occurs (see GBFLWupdate2022Extras.pdf available at SASINF for discussion of sensitivity testing on this and other potential issues).

- If the stock status has changed a lot since the previous assessment, explain why this occurred.

The stock status of Georges Bank Winter Flounder has changed from 'overfished and overfishing is not occurring' to 'not overfished and overfishing is not occurring'.

- Provide qualitative statements describing the condition of the stock that relate to stock status.
The 'Plan B' assessment results (available at [SASINF](#)) indicate that biomass has increased since 2019. There are indications of improvement in stock condition. Catch weight-at-age has been increasing for the last few years and there are indications of a better than average recruitment class in 2020 in the [CA](#) spring survey.
- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.
The Georges Bank Winter Flounder assessment could be improved with a shift to a model that incorporates statistical fits to commercial length and age composition and deprecates the requirement that catch be known without error.
- Are there other important issues?
2020 commercial data, in addition to survey data, was likely affected by the [Covid-19](#) outbreak. Commercial vessels may have carried fewer observers and fished fewer days. The lack of consistency in commercial data may reduce the precision and accuracy of the Georges Bank Winter Flounder assessment in the near term.



Pseudopleuronectes americanus, Winter Flounder.

4.1. Reviewer Comments: Georges Bank Winter Flounder

The 2022 assessment of Georges Bank winter flounder (*Pseudopleuronectes americanus*) updates the 2020 operational **adapt-VPA** assessment that included data for 1982–2019 (NEFSC 2020)¹². This assessment updates commercial fishery catch data, research survey biomass indices, and the analytical **VPA** assessment model and reference points through 2021. Additionally, stock projections have been updated through 2025. Note that the stock is in a revised rebuilding plan, based on fishing at 70% of $F_{MSY\ proxy}$, with rebuilding by 2029.

Unadjusted spawning stock biomass (SSB) in 2021 was estimated to be 7,159 **mt** and the 2021 fully selected fishing mortality (F) was estimated to be 0.049. However, the 2021 point estimate of SSB and F , when adjusted for retrospective error (0.59% for SSB and -0.36% for F), are outside the 90% confidence intervals of the unadjusted 2021 point estimates. Therefore, the values used in the stock status determination were the retrospective-adjusted values of $F_{2021} = 0.076$ which is 17% of the 2022 overfishing threshold ($F_{MSY\ proxy} = 0.452$), and $SSB_{2021} = 4,503\text{ mt}$ which is 60% of the biomass target for an overfished stock (2022 $SSB_{MSY\ proxy} = 7,503$ with a threshold of 50% of $SSB_{MSY\ proxy}$).

Based on this updated assessment, the Georges Bank winter flounder stock is not overfished and overfishing is not occurring.

Short-term projections of biomass were derived by sampling from a cumulative distribution function of recruitment estimates (1982–2020 year class) from the final run of the **adapt-VPA** model. The annual fishery selectivity, maturity ogive (a 3-year moving window), and mean weights-at-age used in the projection are the most recent five-year averages (2017–2021). An SSB retrospective adjustment factor of 0.629 was applied in the projections.

The estimated catch for 2022 is 278 **mt**, which results in catch advice of 2360, 1963 and 1819 **mt** for 2023–2025 respectively.

*Though the estimates of retrospective pattern have declined (at least for SSB), the Panel noted the persistence of retrospective bias in this assessment. This could be an artifact of the **VPA** model, and an evaluation of retrospective patterns in the winter flounder stocks could be insightful. Ultimately, the Panel could see no utility in going to an index-based approach and recommended the **VPA** model be accepted as is.*

Recommendations:

- *Consider using change point analysis to identify recruitment stanzas.*
- *Consider statistical catch at age model in next Research Track.*

*The Panel concluded that the 2022 assessment update for Georges Bank winter flounder fulfilled the recommendations of the **AOP**, is technically sufficient to evaluate stock status and provide scientific advice and meets the Terms of Reference for the stock's assessment. The assessment represents Best Scientific Information Available (**BSIA**) for this stock for management purposes.*

¹²NEFSC. 2022. Operational Assessment of 14 Northeast Groundfish Stocks Updated Through 2018. US Dep Commer, NOAA Fisheries, Northeast Fish Sci Cent Ref Doc. 22-06; 227p. [CRD22-06](#)

References:

Miller, T.J., Richardson, D., Politis, P., Blaylock, J., Manderson, J. and Roebuck, C. 2020. Relative efficiency of a chain sweep and the rockhopper sweep used for the NEFSC bottom trawl survey and biomass estimates for winter and windowpane flounder and red hake stocks. In press. US Dept. Commer., Northeast Fish. Sci. Cent. Ref. Doc. 20-XX; 31p. [CRD20-XX](#)

Northeast Fisheries Science Center. 2020. Operational assessment of 14 Northeast groundfish stocks, updated through 2019. In press. US Dept. Commer., Northeast Fish. Sci. Cent. Ref. Doc. 20-XX; 212p. [CRD20-XX](#)



Displaying a Winter flounder catch.




Figure 10: Trends in spawning stock biomass (mt) of Georges Bank Winter Flounder between 1982 and 2021 from the current (solid line) and previous (dashed line) assessments and the corresponding $SSB_{Threshold}$ ($\frac{1}{2}SSB_{MSY proxy}$; horizontal dashed line) as well as SSB_{Target} ($SSB_{MSY proxy}$; horizontal dotted line) based on the 2022 assessment. Biomass was adjusted for a retrospective pattern and the adjustment is shown in red. The 90% normal confidence interval is shown for 2021.




Figure 11: Trends in fully selected fishing mortality (F_{Full}) of Georges Bank Winter Flounder between 1982 and 2021 from the current (solid line) and previous (dashed line) assessments and the corresponding $F_{Threshold}$ ($F_{MSY proxy} = 0.452$; horizontal dashed line) as well as ($F_{Target} = 75\%$ of $F_{MSY proxy}$; horizontal dotted line). F_{Full} was adjusted for a retrospective pattern and the adjustment is shown in red. The 90% normal confidence interval is shown for 2021.




Figure 12: Trends in Recruits (age-1) (000s) of Georges Bank Winter Flounder between 1982 and 2021 from the current (solid line) and previous (dashed line) assessments.




Figure 13: Total catches (mt) of Georges Bank Winter Flounder between 1982 and 2022 by country and disposition (landings and discards).




Figure 14: Indices of abundance for the Georges Bank Winter Flounder for the Northeast Fisheries Science Center (NEFSC) spring (1968–2021) and fall (1963–2021) bottom trawl surveys and the Canadian DFO spring survey (1987–2021). The 90% normal confidence interval is shown.

5. GEORGES BANK HADDOCK

Liz Brooks

*This assessment of the Georges Bank haddock (*Melanogrammus aeglefinus*) stock is a Level 2 Management Track assessment of the 2021 research track assessment, which used the WHAM framework. Prior to the 2021 research track, the last benchmark for this stock was in 2008 (Brooks et al., 2008). Based on the previous update assessment in 2019 (NEFSC, 2022), the stock was not overfished, and overfishing was not occurring. Stock status was not reported for the 2021 research track, but the stock was not overfished and overfishing was not occurring. This assessment updates commercial fishery catch data, research survey indices of abundance, weights and maturity at age, and the WHAM assessment model and reference points through 2021. Stock projections have been updated through 2025. This report reflects decisions made during the Peer Review September 19–22, 2022.*

State of Stock: Based on this updated assessment, the Georges Bank haddock (*Melanogrammus aeglefinus*) stock is not overfished, and overfishing is not occurring (Figures 15–16). Retrospective adjustments were not made to the model results. Spawning stock biomass (*SSB*) in 2021 was estimated to be 79,513 mt which is 66% of the biomass target ($SSB_{MSY\ proxy} = 120,580$; Figure 15). The 2021 average fishing mortality on ages 5–7 was estimated to be 0.137 which is 55% of the overfishing threshold proxy ($F_{MSY\ proxy} = 0.25$; Figure 16). The $F_{MSY\ proxy}$ is expressed as the average F on ages 5–7.

Table 10: Catch and status table for Georges Bank haddock. All weights are in (mt), recruitment is in (000s), and $\bar{F}_{5:7}$ is the average fishing mortality on ages 5 to 7. Model results are from the current updated WHAM assessment. A ρ -adjustment was not applied to values in this Table.

	2014	2015	2016	2017	2018	2019	2020	2021
	<i>Data</i>							
US Commercial discards	1,409	1,552	1,880	786	410	306	178	49
US Commercial landings	4,240	4,762	3,682	3,217	4,017	5,252	6,648	3,641
Canadian Catch	12,953	14,374	11,713	13,384	12,222	14,160	11,052	7,001
Catch for Assessment	18,601	20,687	17,274	17,387	16,647	19,719	17,878	10,691
	<i>Model Results</i>							
Spawning Stock Biomass	130,266	182,309	187,864	218,393	131,917	114,415	99,365	79,513
$\bar{F}_{5:7}$	0.377	0.313	0.208	0.148	0.146	0.18	0.198	0.137
Recruits (age-1)	1,504,138	67,659	109,014	114,816	20,336	28,142	16,332	96,151

Table 11: Comparison of reference points estimated from the 2019 VPA assessment and from the current assessment update. An $F_{40\%SPR}$ proxy was used for the overfishing threshold. The medians and 95% probability intervals are reported for MSY , SSB_{MSY} , and R_{MSY} , based on WHAM projections with fishing mortality fixed at $F_{40\%SPR}$.

	2019	2022
$F_{MSY\ proxy}$	0.33	0.25
SSB_{MSY} (mt)	138,924	120,580 (94,687–153,555)
MSY (mt)	24,400	25,494 (19,979–32,533)
Median recruits (age-1) (000s)	59,143	25,607 (835–785,516)
Overfishing	No	No
Overfished	No	No

Projections: Short term projections were conducted in WHAM, which propagates uncertainty in the processes of recruitment and transitions between numbers at age. For projection specifications, the Plan Development Team supplied an estimate of total catch for 2022, and fishing mortality was set equal to $F_{40\%SPR}$ for 2023–2025. Annual fishery selectivity and maturity were fixed at a recent 2-year average (2020–2021 values), following analyses and decisions made at the 2021 research track. Weights at age for catch and SSB that were predicted from a Gaussian Markov Random Field (GMRF) model, rather than a recent 2-year average, were preferred by the peer reviewers for this Management Track, and were used in the projections summarized in this report. Retrospective adjustments were not applied. The Overfished threshold is 60,290 mt, and the stock is not projected to drop below this value in 2025.

Table 12: Short term projections of total fishery catch and spawning stock biomass (with 95% CI) for Georges Bank haddock based on a harvest scenario of fishing at 100% $F_{MSY\ proxy}$ between 2023 and 2025. Catch in 2022 was assumed to be 9,914 mt (estimate provided by the Groundfish Plan Development Team).

Year	Catch (mt)	SSB (mt)	$\bar{F}_{5:7}$
2022	9,914	79,457 (39,624–159,332)	0.147 (0.069–0.311)
Year	Catch (mt)	SSB (mt)	$\bar{F}_{5:7}$
2023	18,482 (7,332–46,591)	90,073 (35,695–227,286)	0.272
2024	17,287 (5,680–52,616)	81,027 (25,060–261,981)	0.272
2025	14,555 (3,926–53,958)	69,916 (17,543–278,641)	0.272

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

Sources of uncertainty include dynamics in the plus group, the magnitude of the 2020 and 2021 year classes, and future assumptions about weights and selectivity at age. The 2013 year class, the largest ever observed for this stock, accounts for 20% of the population abundance in 2021 (at age-8), and is in the plus group for all of the projections. It's contribution to catch (in biomass) in the projections is 35% in 2022, and diminishes to 17% in 2025. However, negative annual deviations have been estimated in the plus group in recent years, and it is uncertain if this will persist in projections. Catches in 2023–2025 are reliant on the 2020 year class, which constitutes 33%–41% of the 2023–2025 catch (in biomass). The 2022 year class is not part of the model input, but initial observations in the survey suggest that it may be close to the time series average; additional observations in future surveys are needed to confirm this. Projections from the research track assessment (with data through 2019) aligned well with estimates from the current assessment model (updated with data through 2021), in spite of the projected selectivities being consistently less than the model estimated selectivities from the current assessment. The accuracy of projected weights varied based on the year and year class, with some being very accurate and others over- or underestimated. A sensitivity projection was made using weights estimated from a Gaussian Markov random field (**GMRF**, methodology in Nielsen, manuscript in preparation), and the review panel recommended using these for projections instead of the 2-year average weights at age. These **GMRF** weights at age predicted a slightly greater increase in weights at age in later years of projections (with large uncertainty bounds), and consequently produced larger estimates of catch and **SSB** in 2022–2025 compared to projections using a two year average for weights at age. Long range accuracy for projecting weights and selectivity is not expected, given the many factors that influence those processes.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted **SSB** or $\bar{F}_{5;7}$ lies outside of the approximate joint confidence region for **SSB** and $\bar{F}_{5;7}$).

The 7-year Mohn's ρ , relative to **SSB**, was 0.70 in the 2019 assessment and was 0.26 in 2021. The 7-year Mohn's ρ , relative to **F**, was –0.44 in the 2019 assessment and was –0.27 in 2021. There was a minor retrospective pattern for this assessment because the ρ -adjusted estimates of 2021 **SSB** ($SSB_{\rho} = 79,513$) and 2021 **F** ($F_{\rho} = 0.137$) were inside the estimated 95% confidence regions around **SSB** (46,084–137,174) and **F** (0.073–0.259). No retrospective adjustment was made for either the determination of stock status or for projections of catch in 2023.

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?

As noted in the first bullet, population projections for Georges Bank haddock are uncertain due to future values of selectivity and weights at age, dynamics of the plus group, and magnitude of incoming 2020 and 2021 year classes. This stock is not in a rebuilding plan.

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the affect these changes had on the assessment and stock status.

No changes, other than the incorporation of new data, were made to the Georges Bank haddock assessment for this update. **NEFSC** indices from 2009–2021 were calculated using tow-specific swept area.

- If the stock status has changed a lot since the previous assessment, explain why this occurred.

The stock status of Georges Bank haddock has not changed.

- Provide qualitative statements describing the condition of the stock that relate to stock status.


*The Georges Bank haddock shows a broad age structure, and broad spatial distribution. This stock has produced several exceptionally strong year classes in the last 20 years, leading to record high **SSB** in the last decade. As the strong year classes age out of the population, abundance has returned to levels last observed in the early 2000s, which could potentially lead to an increase in weights at age as growth is released from density-dependent pressures. Catches in recent years have been well below the total quota (US+Canada), but projected catch levels will be substantially less than recent quotas due to declining abundance and the combined effect of re-estimated Canadian weights at age and a re-estimated length-based calibration for the **NEFSC Albatross** : **H.B. Bigelow** vessels.*

- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.

The research track assessment in 2021 for Georges Bank haddock strongly recommended studies to collect data to re-estimate gutted to whole weight conversion factors, as well as measuring individual fish weight in addition to the length and otolith sampling performed on commercially landed fish.

- Are there other important issues?

*The Georges Bank haddock assessment estimates that the haddock stock has declined to levels last observed in the early 2000s. Projections at $F_{40\%SPR}$ using **GMRF** weights at age predict a slight increase in **SSB** in 2023 but then a decline in 2024–2025. Projections at $F_{40\%SPR}$ using a 2-year average for weights-at-age predicted a steady decline in **SSB**, with the stock on the cusp of overfished in 2024, and overfished in 2025. Future stock status is very dependent on assumed weights at age. Surges in stock abundance and quotas are driven by strong year classes, creating a boom and bust cycle. The current assessment shows the stock leaving the boom phase and heading in the bust direction. If initial estimates of the 2020 and 2021 year classes are at or above the time series average, this may slow the current decline.*



Melanogrammus aeglefinus, Haddock.

5.1. Reviewer Comments: Georges Bank haddock

The 2022 assessment of the Georges Bank haddock (*Melanogrammus aeglefinus*) stock is the Management Track update of the 2022 Research Track assessment and also of the 2019 Operational Assessment (NEFSC 2022)¹³. Note that this assessment was also peer reviewed as part of the 2022 haddock Research Track assessment. Prior to the 2021–2022 research track, the last benchmark assessment for this stock was in 2008 (Brooks et al., 2008)¹⁴.

This assessment utilizes the Woods Hole Assessment Model (WHAM) and updates commercial fishery catch data, research survey indices of abundance, weights and maturity at age, and reference points through 2021. Stock projections have been updated through 2025.

Retrospective adjustments of the model results were not necessary. Spawning stock biomass (*SSB*) in 2021 was estimated to be 79,513 mt which is 66% of the biomass target ($SSB_{MSY\ proxy} = 120,580\ mt$). The 2021 average fishing mortality on ages 5–7 was estimated to be 0.137 which is 55% of the overfishing threshold proxy ($F_{MSY\ proxy} = 0.25$). The *F* is expressed as the average *F* on ages 5–7, and $F_{MSY\ proxy}$ is $F_{40\%SPR}$.

Stock status was not reported for the 2021–2022 research track but based on this analysis the stock was not overfished and overfishing was not occurring.

Short term projections were conducted in WHAM, which propagates uncertainty in the processes of recruitment and deviations in transitions between ages. For projection specifications, the Plan Development Team supplied an estimate of total catch for 2022, and then fishing mortality was set equal to $F_{40\%SPR}$ for 2023–2025. Annual fishery selectivity, maturity, and weights at age were fixed at 2020–2021 values, following analyses and decisions made at the 2021 research track. Weights at age for catch and *SSB* that were predicted from a Gaussian Markov Random Field (GMRF) model, rather than a recent 2-year average, were preferred by the peer reviewers for this management track and were used in the projections summarized in this report. The overfished threshold is 60,290 mt, and the stock is not projected to drop below this value in 2025.

The estimated catch for 2022 is 9914 mt, which results in catch advice of 18482, 17287, and 14555 mt for 2023–2025, respectively.

The Panel's discussion on this assessment focused on the choice of Weight At Age (WAA) and selectivity functions to use in the projections. The Panel noted that the two functions seemed to show opposite trajectories over the past decade: WAA was decreasing while selectivity at age was increasing, which suggests that size-based selectivity is possible. However, the latter could have also been affected by fleet targeting and management changes.

¹³NEFSC. 2022. Operational Assessment of 14 Northeast Groundfish Stocks Updated Through 2018. US Dep Commer, NOAA Fisheries, Northeast Fish Sci Cent Ref Doc. 22-06; 227p. [CRD22-06](#)

¹⁴Brooks, E.N, M.L. Traver, S.J. Sutherland, L. Van Eeckhaute, and L. Col. 2008. In Northeast Fisheries Science Center. 2008. Assessment of 19 Northeast Groundfish Stocks through 2007: Report of the 3rd Groundfish Assessment Review Meeting (GARM III), Northeast Fisheries Science Center, Woods Hole, Massachusetts, August 4–8, 2008. US Dep Commer, NOAA Fisheries, Northeast Fish Sci Cent Ref Doc. 08-15; 884p+xvii. [CRD08-15](#)

Nonetheless, the question remained about which of the two approaches to **WAA** (i.e., two-year averaging or Gaussian Markov Random Field model [**GMRF**]) should be used for projections. Ultimately, the Panel recommended that the **GMRF** approach be used, in part because of its quick response to changes and because a model-based approach was more robust than a two-year average. The Panel recommends that the analyst continue to evaluate the utility/accuracy of **GMRF** vs averaging.


Finally, the Panel acknowledged that the eastern Georges Bank haddock stock assessment, also **WHAM** based, has produced a different interpretation of stock's status and trajectory.

The Panel concluded that the 2022 assessment update for Georges Bank haddock fulfilled the recommendations of the **AOP**, is technically sufficient to evaluate stock status and provide scientific advice and meets the Terms of Reference for the stock's assessment. The assessment represents Best Scientific Information Available (**BSIA**) for this stock for management purposes.

References:

Brooks, E.N, M.L. Traver, S.J. Sutherland, L. Van Eeckhaute, and L. Col. 2008. In Northeast Fisheries Science Center, 2008. Assessment of 19 Northeast Groundfish Stocks through 2007: Report of the 3rd Groundfish Assessment Review Meeting (**GARM III**), Northeast Fisheries Science Center, Woods Hole, Massachusetts, August 4–8, 2008. US Dept Commer, **NOAA** Fisheries, Northeast Fish Sci Cent Ref Doc. 08-15; 884p.+xvii. [CRD08-15](#)

Northeast Fisheries Science Center. 2022. Stock Assessment Update of 14 Northeast Groundfish Stocks Through 2018. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 22-06; 232p. [CRD22-06](#)



Haddock swimming over rocky bottom.




Figure 15: Trends in spawning stock biomass of Georges Bank haddock between 1931 and 2021 from the current (solid line) and previous (dashed line) 2019 assessment and the corresponding $SSB_{Threshold}$ ($\frac{1}{2}SSB_{MSY proxy}$; horizontal dashed line) as well as SSB_{Target} ($SSB_{MSY proxy}$; horizontal dotted line) based on the 2021 assessment. The 95% confidence intervals are shown.




Figure 16: Trends in the average fishing mortality ($\bar{F}_{5:7}$) of Georges Bank haddock between 1931 and 2021 from the current (solid line) and previous (dashed line) assessment and corresponding $F_{\text{Threshold}}$ ($F_{\text{MSY proxy}} = 0.25$; horizontal dashed line) based on the 2021 assessment. $\bar{F}_{5:7}$ was adjusted for a retrospective pattern and the adjustment is shown in red. The 95% confidence intervals are shown.




Figure 17: Trends in Recruits (age-1) (000s) of Georges Bank haddock between 1931 and 2021 from the current (solid line) and previous (dashed line) assessment. The 95% confidence intervals are shown.




Figure 18: Total catch of Georges Bank haddock between 1931 and 2021 by fleet (US Commercial, Canadian, or foreign fleet) and disposition (landings and discards).




Figure 19: Indices of biomass (Mean kg/tow) for the Georges Bank haddock stock between 1963 and 2021 for the Northeast Fisheries Science Center (NEFSC) spring and fall bottom trawl surveys and the DFO winter bottom trawl survey. The approximate 95% log-normal confidence intervals are shown for DFO only. Confidence bounds for the new length-based biomass calibration are not yet available.

6. GULF OF MAINE HADDOCK

Charles Perretti

*This assessment of the Gulf of Maine haddock (*Melanogrammus aeglefinus*) stock is a Management Track update assessment of the existing benchmark assessment (NEFSC, In Prep). Based on the previous Management Track assessment (NEFSC 2022), the stock was not overfished, and overfishing was not occurring. This assessment updates commercial and recreational fishery catch data, research survey indices of abundance, and the analytical ASAP assessment model and reference points through 2021. Additionally, stock projections have been updated through 2025*

State of Stock: Based on this updated assessment, the stock status for the Gulf of Maine haddock (*Melanogrammus aeglefinus*) stock is not overfished and overfishing is occurring (Figures 20–21). Retrospective adjustments were not made to the model results (see **Special Comments** section of this report). Spawning stock biomass (*SSB*) in 2021 was estimated to be 16,528 (mt) which is 270% of the biomass target ($SSB_{MSY\ proxy} = 6,123$; Figure 20). The 2021 fully selected fishing mortality was estimated to be 0.375 which is 111% of the overfishing threshold proxy ($F_{MSY\ proxy} = F_{40\%SPR} = 0.338$; Figure 21).

Table 13: Catch and status table for Gulf of Maine haddock. All weights are in (mt) recruitment is in (000s) and F_{Full} is the fully selected fishing mortality. Model results are from the current updated ASAP assessment.

	2014	2015	2016	2017	2018	2019	2020	2021
	<i>Data</i>							
Recreational discards	617	526	966	743	312	234	271	137
Recreational landings	456	295	1,026	1,787	801	778	909	722
Commercial discards	22	42	72	91	54	66	122	25
Commercial landings	314	650	1,342	2,273	2,542	3,464	3,556	3,718
Foreign landings	0	0	0	0	0	0	0	0
Catch for Assessment	1,410	1,512	3,406	4,894	3,710	4,542	4,858	4,602
	<i>Model Results</i>							
Spawning Stock Biomass	9,787	19,437	32,445	36,189	34,312	29,887	22,988	16,528
F_{Full}	0.346	0.217	0.278	0.27	0.176	0.212	0.282	0.375
Recruits (age-1)	83,932	4,915	5,169	6,976	5,629	2,080	3,059	22,781

Projections: Short term projections of median total fishery yield and spawning stock biomass for Gulf of Maine haddock were conducted based on a harvest scenario of fishing at the $F_{MSY\ proxy}$ between 2023 and 2025. Catch in 2022 has been estimated at 3,912 mt. Recruitment was sampled from a cumulative distribution function of model estimated age-1 recruitment from 1977–2019. The age-1 estimate in 2022 was generated from the geometric mean of the 1977–2021 recruitment series. The annual fishery selectivity in the projections was the terminal selectivity from the assessment model. The time-invariant

maturity ogive, and the projected mean weights-at-age were used in the projections. Retrospective adjustments were not applied in the projections.

Table 14: Comparison of reference points estimated in an earlier assessment and from the current operational assessment. The overfishing threshold is the $F_{MSY proxy}$ ($F_{40\%SPR}$). The biomass target, ($SSB_{MSY proxy}$) was based on long-term stochastic projections of fishing at the $F_{MSY proxy}$. Median recruitment reflects the median estimated age-1 recruitment from 1977–2019. Intervals shown reflect the 5th and 95th percentiles.

	2019	2022
$F_{MSY proxy}$	0.341 (0.281–0.412)	0.338 (0.254–0.458)
SSB_{MSY} (mt)	7,993 (3,218–34,191)	6,123 (2,864–19,628)
MSY (mt)	1,597 (651–6,797)	1,098 (507–3,533)
Median recruits (age-1) (000s)	1,789 (285–17,883)	1,753 (283–12,832)
Overfishing	No	Yes
Overfished	No	No

Table 15: Short term projections of total fishery catch and spawning stock biomass for Gulf of Maine haddock based on a harvest scenario of fishing at $F_{MSY proxy}$ ($F_{40\%SPR}$) between 2023 and 2025. Catch in 2022 was assumed to be 3,912 (mt).

Year	Catch (mt)	SSB (mt)	F_{Full}
2022	3,912	14,920	0.471
Year	Catch (mt)	SSB (mt)	F_{Full}
2023	2,515	16,002	0.338
2024	2,552	15,400	0.338
2025	2,434	13,189	0.338

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

The main source of uncertainty in this assessment is the faster than expected rate of decline in biomass that has occurred since the last update. Although the last update projected a decline in biomass, the realized decline has been faster than anticipated, and this has resulted in a substantial increase in F which is now above the overfishing threshold.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted SSB or F_{Full} lie outside of the approximate joint confidence region for SSB and F_{Full}).

This assessment exhibits a retrospective pattern, however the 7-year Mohn’s ρ -adjusted values of SSB and F fall within the 90% confidence interval of their estimates, therefore the pattern is considered minor, and no retrospective adjustment is required.

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?
Population projections for Gulf of Maine haddock in the previous update correctly identified that the population would decline, however the rate of decline has been faster than projected. As a result, the projected 2021 SSB from the last assessment is above the upper confidence bound of the 2021 SSB estimated in the current assessment. The over-projection of SSB was magnified by a change in direction of the retrospective pattern and over-projected weights-at-age, the latter of which has been addressed in this update using a new weight-at-age projection model. The projected F for 2022 is above the overfishing threshold (Table 15). This stock is not in a rebuilding plan.
- Describe any changes that were made to the current stock assessment beyond incorporating additional years of data, and the effect these changes had on the assessment and stock status.
The actual tow-swept-area was used when calculating the NEFSC Bottom Trawl indices of abundance for the R/V Bigelow years (2009+). A comparison of the survey time series with and without this adjustment showed that the impact on the time series is minor. In addition, following the recommendation of the 2021 Research Track Review Panel (CIE, In Prep), an exploratory model run was performed which included the NEFSC Bottom Longline Survey (BLLS). Model diagnostics were generally worse when including the BLLS, particularly the retrospective pattern, and therefore the Base model was chosen as the preferred model. Also following the recommendation of the Research Track Review Panel, a state-space growth model was developed for projecting WAA. In a 20-year retrospective backtest, the growth model outperformed the previous approach of using the terminal two-year average, and therefore it was chosen as the WAA projection method in this update.
- If the stock status has changed a lot since the previous assessment, explain why this occurred.
Unlike the previous assessment (NEFSC 2022), overfishing is now occurring. See above comments on why this occurred. The stock remains not overfished.
- Provide qualitative statements describing the condition of the stock that relate to stock status.
The Gulf of Maine haddock stock has experienced several large year classes since 2010, particularly the 2013 year class. The stock has recently declined as these large year classes have aged out. The 2020 year class is currently estimated as the second largest on record, however it is still substantially smaller than the 2013 year class and its estimate is highly uncertain. Future stock status will depend on the strength of this and subsequent year classes.
- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.
A better understanding of what is driving the retrospective pattern would be helpful.
- Are there other important issues?
None.

6.1. Reviewer Comments: Gulf of Maine haddock

The 2022 assessment of the Gulf of Maine haddock (*Melanogrammus aeglefinus*) stock is the Management Track update of the 2021 Research Track assessment and 2019 Operational Assessment (NEFSC 2022)¹⁵. Note that this assessment was also peer reviewed as part of the 2021 haddock Research Track assessment. This assessment updates commercial and recreational fishery catch data, research survey indices of abundance, and the analytical ASAP assessment model and reference points through 2021. Stock projections have been updated through 2025.

The Panel concluded that the 2022 assessment update for Gulf of Maine haddock fulfilled the recommendations of the AOP, is technically sufficient to evaluate stock status and provide scientific advice and meets the Terms of Reference for the stock's assessment. The assessment represents Best Scientific Information Available (BSIA) for this stock for management purposes.

Terms of Reference: Gulf of Maine haddock

1. Estimate catch from all sources including landings and discards.

Commercial and recreational discards, landings, and age composition were updated through 2021.

An issue with the commercial Numbers and Weight-At-Age (NAA/WAA) exists for 2021 in that only one 'Large' fish was sampled in 2021. Only 6% of the catch was Large, but the effect of this low sampling was investigated further by considering three solutions: 1) Assign all Large catch to the one observed length, 2) borrow the 2020 Large samples and expand them annually, and 3) borrow the 2020 Large samples and expand them quarterly. After examining NAA and its CV, and WAA, the analyst determined that use of the 2020 Large samples with quarterly expansion provided the most robust solution.

The Panel was concerned about the low level of Port side sampling of commercial landings that has occurred since 2019.

2. Evaluate indices used in the assessment (e.g., indices of relative or absolute abundance, recruitment, state surveys, age-length data, etc.).

This TOR was satisfactorily addressed.

Spring and fall tow-specific swept area NEFSC bottom trawl survey indices as well as survey maturities, and weights at age were updated through 2021 (2020 surveys were not conducted). The NEFSC 2014–2021 Bottom Longline Survey results were also considered for inclusion in the model.

3. Estimate annual fishing mortality, recruitment and stock biomass (both total and spawning stock) as possible (depending on the assessment method) for the time series using the approved assessment method and estimate their uncertainty. Include retrospective analyses if possible (both historical and within-model) to allow a comparison with previous assessment results and projections, and to examine model fit.

¹⁵NEFSC 2022. Operational Assessment of 14 Northeast Groundfish Stocks Updated Through 2018. US Dep Commer, NOAA Fisheries, Northeast Fish Sci Cent Ref Doc. 22-06; 227p. [CRD22-06](#)

- (a) Include bridge runs to sequentially document each change from the previously accepted model to the updated model proposed for this peer review.
- (b) Prepare a backup assessment approach that would serve as an alternative for providing scientific advice to management if the analytical assessment were to not pass review.

The Base model used here was the final accepted **ASAP** model from the 2021 Haddock Research Track Assessment for 1977–2021 (1977 is the first year catch-at-age is available). Maturity was from the 1977–2021 time series average, assumed natural mortality $M = 0.2$, and **SSB** weights-at-age use Rivard calculation. Commercial and recreational catches were for ages 1–9+ with Weights-at-Age (**WAA**) estimated from the most recent 5-year average. A combined commercial and recreational fleet was modeled with three selectivity blocks (at-age) 1977–1988, 1989–2004, 2005+. For the surveys, selectivity was freely estimated at-age (fixed 4+ (spring), 6+ (fall)) and catchability was fixed over time.

The **NEFSC** Bottom Longline Survey (**BLLS**) was included as an exploratory model in addition to the **NEFSC** Bottom Trawl Survey data, as recommended by the **GOM** Haddock Research Track Review Panel. The configuration of the **BLLS** model was identical to the Base model, with the addition of the **BLLS** survey. **BLLS** selectivity was freely estimated at-age and fixed for ages 6+ (spring) and 5+ (fall).

A Bridge run was prepared to compare the 2019 Management Track (**MT**) model with the 2021 Research Track (**RT**) model with the updated 2022 **MT** original base model and base+**BLLS**. Model estimates of **SSB**, F , and recruits were comparable.

*The Panel spent significant time discussing whether to include the **BLLS** in this assessment. While the Panel agreed the **BLLS** provided additional useful information with the model providing a good fit for the **BLLS** survey results, overall diagnostics suggested it degraded the quality of the model. It increased Mohn's ρ from 0.30 in the 2022 base model to 0.48 in the **BLLS** model, which was a doubling of the ρ from the 2021 **RT** model. Ultimately, the Panel agreed to recommend the **BLLS** not be included in the 2022 **MT** model but be reconsidered in the next assessment once 2 more years of data are available.*

- 4. Re-estimate or update the **BRPs** as defined by the management track level and recommend stock status. Also, provide qualitative descriptions of stock status based on simple indicators/metrics (e.g., age- and size-structure, temporal trends in population size or recruitment indices, etc.).

This **TOR** was satisfactorily addressed.

Retrospective adjustments were not made to the model. Spawning stock biomass (**SSB**) in 2021 was estimated to be 16,528 **mt** which is 270% of the biomass target ($SSB_{MSY\ proxy} = 6,123\ \text{mt}$). The 2021 fully selected fishing mortality was estimated to be 0.375 which is 111% of the overfishing threshold proxy ($F_{MSY\ proxy} = F_{40\%SPR} = 0.338$).

Based on this updated assessment, the stock's status is not overfished and overfishing is occurring.

- 5. Conduct short-term stock projections when appropriate.

This **TOR** was satisfactorily addressed.

Short term projections of median total fishery yield and spawning stock biomass for Gulf of Maine haddock were conducted based on a harvest scenario of fishing at the F_{MSY} proxy between 2023 and 2025. Recruitment was sampled from a cumulative distribution function of model estimated age-1 recruitment from 1977–2019. The age-1 estimate in 2022 was generated from the geometric mean of the 1977–2021 recruitment series. The annual fishery selectivity in the projections was the terminal selectivity from the assessment model. The time-invariant maturity ogive and projected mean weights at age derived from a state-space growth model were used in the projections. Retrospective adjustments were not applied in the projections.

The estimated catch for 2022 is 3912 **mt**, which results in catch advice of 2515, 2552, and 2434 **mt** for 2023–2025, respectively.

*The Panel discussed whether to use the recent 2-year average or the state-space growth model predictions for estimates of **WAA** in projections. The growth model tracks growth curves of cohorts to predict future **WAA** and results in lower projections of **WAA** in the future than the 2-year average with lower projected catch in 2023–2024 and higher **SSB**. Based on the growth model fits to catch weights and **SSB** weights, the Panel recommended using the growth model for **WAA** in the projections.*

6. Respond to any review panel comments or **SSC** concerns from the most recent prior research or management track assessment.

This **TOR** was partially addressed in that the Research Track Peer Review Panel recommendations on the Bottom Longline Survey and the Weight-At-Age analysis were fully addressed by the assessor. Other items remaining to be dealt with by the assessor include consideration of variable **M** values to incorporate age and system variability (including the effect on management advice), and the appropriateness of the $F_{40\%SPR}$ proxy. Remaining research from the Review Panel including the generation of testable hypotheses to explain recruitment variability in the Gulf of Maine (including an updating of the fall bloom relationship to recruitment) are not the responsibility of the assessor.

Research recommendations from this Panel included:

- *The **BLLS** should be reconsidered for inclusion in the assessment in the next **MT** assessment, once 2 more years of data are available.*
- *This stock should be considered for evaluation using a state-space framework, which could provide insight into the rapid decline in biomass. The Panel suggested that Gulf of Maine haddock may be included as a case study in the State-Space Modeling Research Track effort to be convened soon.*
- *Low Port sampling of landings is a problem for this and many other stocks. **NMFS** should either return this sampling to pre 2109 levels or consider data from observer sampling.*

References:

Northeast Fisheries Science Center. In prep. Gulf of Maine Haddock Research Track Assessment. US Dept Commer, Northeast Fish Sci Cent Ref Doc. XX-XX. [HaddockWG](#)

Northeast Fisheries Science Center. 2022. Stock Assessment Update of 14 Northeast Groundfish Stocks Through 2018. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 22-06; 232p. [CRD22-06](#)

Center for Independent Experts. In prep. Independent Peer Review Report on the Gulf of Maine Haddock Research Track. [HaddockWG](#)



Sorted Haddock in baskets.




Figure 20: Trends in spawning stock biomass (SSB) of Gulf of Maine haddock between 1977 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $SSB_{\text{Threshold}}$ ($\frac{1}{2}SSB_{MSY \text{ proxy}}$; horizontal dashed line) as well as SSB_{Target} ($SSB_{MSY \text{ proxy}}$; horizontal dotted line) based on the 2022 assessment. The approximate 90% log-normal confidence intervals are shown. The red dot indicates the ρ -adjusted SSB value that would have resulted had a retrospective adjustment been made (see [Special Comments](#) section).




Figure 21: Trends in the fully selected fishing mortality (F) of Gulf of Maine haddock between 1977 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $F_{\text{Threshold}}$ ($F_{\text{MSY proxy}} = 0.338$; horizontal dashed line) from the 2022 assessment model. The approximate 90% log-normal confidence intervals are shown. The red dot indicates the ρ -adjusted F value that would have resulted had a retrospective adjustment been made (see [Special Comments](#) section).




Figure 22: Trends in Recruits (age-1) (000s) of Gulf of Maine haddock between 1977 and 2021 from the current (solid line) and previous (dashed line) assessment. The approximate 90% log-normal confidence intervals are shown.




Figure 23: Total catch of Gulf of Maine haddock between 1977 and 2021 by fleet (commercial, recreational, or foreign) and disposition (landings and discards).




Figure 24: Indices of biomass for the Gulf of Maine haddock between 1963 and 2021 for the Northeast Fisheries Science Center (NEFSC) spring and fall bottom trawl surveys. The approximate 90% log-normal confidence intervals are shown.

7. ATLANTIC HALIBUT

Daniel Hennen

This assessment of the Atlantic halibut (*Hippoglossus hippoglossus*) stock is an update of the existing 2019 ‘Plan B’ assessment (Rago, 2018). This assessment updates commercial fishery catch data, commercial and survey indices of abundance, and the First Second Derivative (FSD) model through 2021. Reference points are unknown and have not been updated.

State of Stock: Based on this updated assessment, Atlantic halibut (*Hippoglossus hippoglossus*) stock status cannot be determined analytically due to a lack of biological reference points associated with the FSD method. Biomass (*SSB*) in 2021 was unknown. The 2021 fully selected fishing mortality was unknown.

Table 16: Catch and status table for Atlantic halibut. All weights are in (mt).

	2014	2015	2016	2017	2018	2019	2020	2021
	<i>Data</i>							
Commercial discards	26	23	31	27	46	75	37	26
Commercial landings	45	62	68	64	54	50	48	39
CA landings	33	30	34	35	46	54	156	120
Catch for Assessment	104	115	133	125	146	178	240	185
	<i>Model Results</i>							
Catch Multiplier	1.02	1.18	1.02	1.02	0.94	0.84	0.92	0.87
Catch Advice	141	106	136	136	128	138	149	220

Table 17: There are no current reference points for Atlantic halibut which is on a ‘Plan B’ assessment that does not allow for the estimation of reference points. Therefore the status of the stock relative to overfishing and overfished status is unknown. Note: based on NOAA policy, the Agency previously decided the stock status was overfished and overfishing not occurring.

	2019	2022
$F_{MSY\ proxy}$	NA	
SSB_{MSY} (mt)	NA	
MSY (mt)	NA	
Overfishing	Unknown	Unknown
Overfished	Unknown	Unknown

Projections: Short term projections are not possible using the FSD approach. The FSD approach is based on applying a multiplier to the catch from the previous year and cannot be projected beyond the catch time series. The catch multiplier for 2021 resulting from the FSD model is 0.87 and the estimated

catch for 2021 is 185 mt, which results in catch advice of 160 mt for 2022. The FSD model is explained in (Rago, 2018) and additional information is available in a document called 'AtlanticHalibutMTextras.pdf', both are available at [SASINF](#).

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

The assessment model (FSD) used for Atlantic halibut is a 'Plan B' assessment method. It uses recent trends in 3 abundance indices as well as recent changes in those trends to adjust the previous year's catch. For example, if the abundance indices are increasing, the catch will be adjusted up. If that increasing trend in abundance is increasing in magnitude over time, the adjustment to catch will be commensurately higher. The FSD method was rigorously tested in simulation (Rago, 2018) and should perform well for Atlantic halibut in the US. Sources of uncertainty in the FSD method include process error related to potential changes in stock productivity over time, the choice of relative weights for the control parameters used in the model and the lag in information inherent in using change in trend as one of the control parameters, which requires dropping one data point from the regression fit to generate a comparison. Other sources of uncertainty include the observation error in the abundance indices. The FSD method also relies on the assumption that abundance can be described with linear dynamics, but that assumption should be relatively unimportant if the stock abundance is well below its theoretical carrying capacity.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major?

The FSD model does not support retrospective analysis.

- Based on this stock assessment, are population projections well determined or uncertain?

The FSD model provides catch advice in the year following the terminal year of the input data. It is not intended to project further ahead than one year. It is possible however to assume that catch in the year following the terminal year will equal the catch advice from the FSD model and that the population abundance indices will continue to follow the same trend and that the change in trend will be identical to the previous five years of data. These assumptions allow for a projection any number of years into the future. The relative quality of these projections degrades as the indices of abundance depart from the behavior of the most recent data available to the model.

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the affect these changes had on the assessment and stock status.

The loss of a survey data point in 2020 (Covid) resulted in the need to interpolate one survey index observation. The survey index value used in 2020 was equal to the average of the 2019 and 2021 values.

Catch efficiency studies and data are not used for the Atlantic halibut assessment because not enough Atlantic halibut are caught to provide a comparison between the gear types and produce estimate of catchability.

- If the stock status has changed a lot since the previous assessment, explain why this occurred.
Stock status cannot be determined and remains unchanged. The stock is likely depleted relative to its virgin biomass based on estimates of historical landings, which were much higher than current landings. Rago in his 2018 report argued that overfishing was unlikely because the catch multiplier estimated in the FSD model had been greater than one for several years. The catch multiplier has now been less than one for four years, which would be consistent with recent overfishing. There is however, no way to determine stock status without reference points.
- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.
The Atlantic halibut assessment could be improved with more precise fishery independent indices of abundance, additional age and length composition data, and a better understanding of stock structure. These would allow for alternative assessment methods, and potential development of a more sophisticated stock assessment model.
- Are there other important issues?
Canadian catch in 2020 and 2021 in area 5Z (eastern Georges Bank) was 4 times higher than it has been in at least the last 20 years (see SASINF). Because this area is included in the calculation of catch in the FSD model, the recommended catch output (catch advice) for 2021 and 2022 is at or above status quo (circa 2017–2019). This result is counter to expectation given that the catch multiplier for each year since 2017 indicates a reduction in recommended catch is warranted (Table 16). Managers should be aware of this issue when recommending catch levels for Atlantic halibut.



Hippoglossus hippoglossus, Atlantic Halibut.

7.1. Reviewer Comments: Atlantic halibut

The 2022 assessment of Atlantic halibut (*Hippoglossus hippoglossus*) updates the 2019 ‘Plan B’ assessment use of the First Second Derivative (FSD) model (Rago 2018)¹⁶ with additional commercial fishery catch data, commercial discard indices and fall bottom trawl survey indices of abundance through 2021.

A ‘Plan B’ assessment does not allow for the estimation of reference points (i.e., F_{MSY} , and SSB_{MSY} cannot be determined).

Short-term projections are not possible using the FSD approach. The FSD approach is based on applying a multiplier to the catch from the previous year and cannot be projected beyond the catch time series. The catch multiplier for 2021 resulting from the FSD model is 0.85 and the estimated catch for 2021 is 174 mt, which results in catch advice of 149 mt for 2022.

The Panel considered the analyst’s recommendation that Atlantic halibut’s stock status should be considered unknown. Given that there is no accepted previous assessment to provide stock status and that the “Plan B” approach does not generate reference points, the Panel strongly recommended listing stock status as unknown.

The Panel raised two concerns with this assessment. The first was the intersection of the US fishery and stock with those of Canada’s fishery and stock. Even though the US assessment tries to account only for the portion of the Atlantic halibut stock that is found on Georges Bank and the Gulf of Maine, it is possible (perhaps likely) that this is only one part of a larger stock of Atlantic halibut that extends onto the Canadian Scotian shelf. Tracking data available from halibut released in the Gulf of Maine suggest a significant number of these fish travel into Canadian waters. The recent high catches of halibut by Canadian fisheries on Georges Bank (e.g., in area 5ZEJ) can only be sustainable if the area is part of a larger stock. As such, the Panel recommended that NMFS and DFO develop a more coordinated approach to the assessment of Atlantic halibut across the region. The Panel endorses at the least a scientific workshop to discuss the stock’s science.

The second, much smaller concern, was how the missing fall 2020 survey was treated in the analysis. Imputing or smoothing by averaging values between the two surveys before and after 2020 was the agreed upon recommendation.

For the next Management Track assessment of the stock, consider different weighting of the commercial discard and survey abundance indices and the potential for use of a CPUE index.

The Panel concluded that the 2022 assessment update for Atlantic halibut fulfilled the recommendations of the AOP and is technically sufficient to provide scientific advice and meets the Terms of Reference for the stock’s assessment. It does not provide sufficient information to evaluate stock status. The assessment represents Best Scientific Information Available (BSIA) for this stock for management purposes.

¹⁶Rago, P.J. 2018. Halibut Assessment Report for 2017 for New England Fishery Management Council, January 24, 2018. Unpublished report. Available online at [SASINF](#).

References:

Rago, P.J. 2018. Halibut Assessment Report for 2017 for New England Fishery Management Council, January 24, 2018. Unpublished, online at [SASINF](#).



Halibut on deck of fishing vessel.




Figure 25: The catch multiplier resulting from the FSD model for Atlantic halibut between 2006 and 2022 from the current (solid line) assessment. A dashed line at 1 is added for reference.




Figure 26: The catch advice resulting from multiplying catch and the catch multiplier from the FSD model for Atlantic halibut between 2006 and 2022 from the current assessment.




Figure 27: Total catch of Atlantic halibut between 2006 and 2022 by disposition (landings and discards).




Figure 28: Indices of biomass for the Atlantic halibut between 2002 and 2021 for the Northeast Fisheries Science Center (NEFSC) fall bottom trawl survey and 2 discard ratio estimators. Discard mortality is assumed to be 0.76 for trawl gear and 0.3 for gillnet gear. The 2020 NEFSC fall bottom trawl value was interpolated as the mean of the 2019 and 2021 values. The 90% lognormal confidence intervals are shown.

8. WHITE HAKE

Katherine Sosebee

This assessment of the white hake (*Urophycis tenuis*) stock is a Management Track update of the 2019 Operational Assessment (NEFSC 2022) and the last benchmark assessment (NEFSC 2013). Based on the previous assessment the stock was overfished and overfishing was not occurring. This assessment updates commercial fishery catch data, research survey indices of biomass, adds an additional survey, and updates the ASAP assessment model and reference points through 2021. Stock projections have been updated through 2025.

State of Stock: Based on this updated assessment, the white hake (*Urophycis tenuis*) stock is not overfished and overfishing is not occurring (Figures 29–30). Retrospective adjustments were made to the model results. Spawning stock biomass (*SSB*) in 2021 was estimated to be 19,497 mt which is 69% of the biomass target ($SSB_{MSY\ proxy} = 28,191$; Figure 29). The 2021 fully selected fishing mortality was estimated to be 0.104 which is 65% of the overfishing threshold proxy ($F_{MSY\ proxy} = 0.1605$; Figure 30).

Table 18: Catch and ASAP results table for white hake. All weights are in (mt) recruitment is in (000s) and F_{Full} is the fishing mortality on fully selected ages (ages 6–9+). Model results are from the current ASAP assessment.

	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
<i>Data</i>										
Commercial discards	50	38	33	24	33	36	29	30	31	32
Commercial landings	2,771	2,235	1,887	1,632	1,325	1,976	1,969	1,975	1,990	1,871
Canadian landings	83	43	35	25	39	32	45	24	83	48
Other landings	0	0	0	0	0	0	0	0	0	0
Catch for Assessment	2,903	2,316	1,955	1,680	1,396	2,043	2,044	2,029	2,104	1,951
<i>Model Results</i>										
Spawning Stock Biomass	21,919	21,867	20,783	19,143	22,186	23,673	19,359	21,276	25,059	23,670
F_{Full}	0.15	0.12	0.1	0.09	0.07	0.09	0.11	0.1	0.09	0.09
Recruits (age-1)	1,991	2,502	2,720	3,082	2,521	2,757	2,975	2,630	1,912	1,740

Projections: Short term projections of catch and *SSB* were derived by sampling from a cumulative distribution function of recruitment estimates from ASAP from 1995–2019. The mean weights-at-age used in the projection are the 2017–2019+2020 averages. The numbers-at-age used to start the projections were adjusted for retrospective bias using age-specific ρ estimates.

Table 19: Comparison of reference points estimated in the 2019 assessment and from the current assessment update. An $F_{40\%SPR}$ proxy was used for the overfishing threshold and SSB_{MSY} was based on long-term stochastic projections which sampled from a cumulative distribution function of recruitment estimates from ASAP from 1963–2019. The annual fishery selectivity, maturity ogive, and mean weights-at-age used in the projection are the most recent 5-year averages.

	2019	2022
$F_{MSY\ proxy}$	0.1677	0.1605
SSB_{MSY} (mt)	31,828	28,191 (22,616–35,424)
MSY (mt)	4,601	4,186 (3,345–5,279)
Median recruits (age-1) (000s)	4,471	4,232
Overfishing	No	No
Overfished	Yes	No

Table 20: Short term projections of total fishery catch and spawning stock biomass for white hake based on a harvest scenario of fishing at $F_{MSY\ proxy}$ between 2023 and 2025. Catch in 2022 was assumed to be 1,964 (mt) which is 56% of the 2022 OFL.

Year	Catch (mt)	SSB (mt)	F_{Full}
2022	1,964	17,978 (15,553–20,679)	0.116
Year	Catch (mt)	SSB (mt)	F_{Full}
2023	2,650	17,679 (15,212–20,216)	0.1605
2024	2,535	17,139 (14,914–19,381)	0.1605
2025	2,547	17,326 (15,360–19,302)	0.1605

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).
 1. *Catch-at-age information is not well characterized due to possible mis-identification of species in the commercial and observer data, particularly in early years, low sampling of commercial landings in some years, and sparse discard length data.*
 2. *Since the commercial catch is aged primarily with survey age/length keys, there is considerable augmentation required, mainly for ages 5 and older. The numbers-at-age and mean weights-at-age in the catch for these ages may therefore not be well specified.*
 3. *White hake may move seasonally into and out of the defined stock area.*
 4. *There are no commercial catch-at-age data prior to 1989 and the catchability of older ages in the surveys is very low. This results in a large uncertainty in starting numbers-at-age.*
 5. *Since 2003, dealers have been culling extra-large fish out of the large category. However, there was no market category for landings until June 2014. The length compositions are distinct from fish characterized as large and have been identified since 2011. This may bias the age composition of the landings, particularly in 2014 when 2000 of the 5000 large samples were these extra-large fish.*

6. A pooled age/length key is used for 1963–1981, fall 2003 survey data as well as the second half of the commercial key, and for the 2020 commercial CAA.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted SSB or F_{Full} lies outside of the approximate joint confidence region for SSB and F_{Full})

The 7-year Mohn's ρ , relative to SSB , was 0.31 in the 2019 assessment and was 0.21 in 2021. The 7-year Mohn's ρ , relative to F , was -0.22 in the 2019 assessment and was -0.17 in 2021. There was a major retrospective pattern for this assessment because the ρ -adjusted estimate of 2021 SSB ($SSB_{\rho} = 19,497$) was outside the approximate 90% confidence regions around SSB (19,894–26,646). A retrospective adjustment was made for both the determination of stock status and for projections of catch in 2023. The retrospective adjustment changed the 2021 SSB from 23,670 to 19,497 and the 2021 F_{Full} from 0.09 to 0.104.

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?

Population projections for white hake are not well determined and projected biomass from the last assessment was near the edge the confidence bounds of the biomass estimated in the current assessment. The rebuilding deadline for this stock is now 2031 and the stock may rebuild.

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the affect these changes had on the assessment and stock status. In the 2022 assessment of white hake, the catch efficiency studies and data were not used because studies were not applicable to roundfish.

Minor changes to the catch-at-age for 2003 and 2015–2018 were made and made little difference to the model. The swept-area adjusted survey values for 2009–2018 were used as well as the bootstrap CVs. This lowered the estimates of SSB over that time period and slightly increased fishing mortality. In addition, one new survey was added to the ASAP model which reduced the retrospective pattern.

- If the stock status has changed a lot since the previous assessment, explain why this occurred.

Stock status of white hake has changed from overfished to not overfished for at least two reasons. First, the retrospective pattern was reduced. Second, the biomass reference point was also reduced because of a lower mean recruitment.

- Provide qualitative statements describing the condition of the stock that relate to stock status.

The white hake stock shows no truncation of age structure. Estimates of commercial landings and discards have decreased over time.

- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.

Age structures collected by the observer program are available and should be aged to augment the survey keys. The additional years of age structures from the ASMFC shrimp survey should also be aged and continue to be collected. There are two bottom longline surveys that should be monitored as the time series gets longer, and the otoliths aged and collected.

- Are there other important issues?

None.

8.1. Reviewer Comments: White hake

The 2022 assessment for white hake (*Urophycis tenuis*) is an update to the 2019 Operational Assessment (NEFSC 2022)¹⁷ and the last benchmark assessment (NEFSC 2013)¹⁸. This assessment updates commercial fishery catch data, research survey indices of biomass, adds additional surveys, and updates the ASAP assessment model and reference points through 2021. Stock projections have been updated through 2025.

The Panel concluded that the 2022 assessment update for white hake fulfilled the recommendations of the AOP, is technically sufficient to evaluate stock status and provide scientific advice and meets the Terms of Reference for the stock's assessment. The assessment represents Best Scientific Information Available (BSIA) for this stock for management purposes.

Terms of Reference: White hake

1. Estimate catch from all sources including landings and discards.

This TOR was satisfactorily addressed.

US and Canadian commercial landings and discard data were updated by market category through 2021 with CAMS tables used for 2020–2021. Catch was split by red/white hake market category using proportion by area fished. Market category of extra-large hake was combined with large hake because of a lack of landings.

US discards were estimated using a method like the NEFSC's Standard Bycatch Reporting Methodology (SBRM) approach by gear type. Longline discards were assumed to be the same Catch-at-Age (CAA) as large-mesh trawl discards, scallop dredge and shrimp trawl discards were the same CAA as small mesh discards and sink gill net discards were the same CAA as landings and discards combined.

The Panel recommends that the SBRM method be used in the next assessment.

2. Evaluate indices used in the assessment (e.g., indices of relative or absolute abundance, recruitment, state surveys, age–length data, etc.).

This TOR was satisfactorily addressed.

Spring and fall tow-specific swept area NEFSC bottom trawl survey indices were updated through 2021 (2020 surveys were not conducted). Additional US survey data were evaluated: the ASMFC shrimp survey (including age and length data) and the NEFSC bottom longline survey. As with other stocks, no 2020 survey data were available except for the fall BLLS. A pooled Age Length Key (ALK) from previous years was used as a proxy for the missing 2020 WAA/CAA.

¹⁷NEFSC. 2022. Operational Assessment of 14 Northeast Groundfish Stocks Updated Through 2018. US Dep Commer, NOAA Fisheries, Northeast Fish Sci Cent Ref Doc. 22-06; 227p. [CRD22-06](#)

¹⁸NEFSC. 2013. 56th Northeast Regional Stock Assessment Workshop (SAW 56) Assessment Report. US Dep Commer, NOAA Fisheries, Northeast Fish Sci Cent Ref Doc. 13-10; 868p. [CRD13-10](#)

3. Estimate annual fishing mortality, recruitment and stock biomass (both total and spawning stock) as possible (depending on the assessment method) for the time series using the approved assessment method and estimate their uncertainty. Include retrospective analyses if possible (both historical and within-model) to allow a comparison with previous assessment results and projections, and to examine model fit.
 - (a) Include bridge runs to sequentially document each change from the previously accepted model to the updated model proposed for this peer review.
 - (b) Prepare a backup assessment approach that would serve as an alternative for providing scientific advice to management if the analytical assessment were to not pass review.

This TOR was satisfactorily addressed.

The Base model used here was the final accepted ASAP model from the preceding Operational Assessment of white hake using catch and survey data for 1963–2021. Catches-At-Age (CAA) for 1963–2018 were updated with additional CAA for 2019–2021. The missing 2020 CAA was considered using pooled Age Length keys (no survey ages) with sensitivity to leaving out 2020 CAA. Survey selectivities were set to fully selected at age 3 and other ages were allowed to be estimated, while fishery selectivity was set to full for ages 6–9. The assessment includes two selectivity blocks 1963–1997 and 1998–2021. Shrimp survey and Bottom Longline Survey indices were both evaluated for incorporation into the model.

Other than the introduction of new data, there were only minor changes to the previous assessment. As a result, bridge runs showed negligible differences from the prior assessment.

The Panel recommended that this assessment include the BTS and the shrimp survey, but not the bottom longline survey (BLLS). This should be reevaluated at the next management track assessment when more data is available for both surveys. They may be useful in that both surveys sample age classes which may not be well sampled by the BTS. Consideration should also be given to including the Maine/New Hampshire surveys in the next assessment.

The Panel also recommended that future assessments consider splitting the Albatross and Bigelow BTS time series.

4. Re-estimate or update the BRPs as defined by the management track level and recommend stock status. Also, provide qualitative descriptions of stock status based on simple indicators/metrics (e.g., age- and size-structure, temporal trends in population size or recruitment indices, etc.).

This TOR was satisfactorily addressed.

The retrospective pattern improved with adjustments made to the model. Spawning stock biomass (SSB) in 2021 was estimated to be 19,369 mt which is 67% of the biomass target ($SSB_{MSY\ proxy} = 28,039$). The 2021 fully selected fishing mortality was estimated to be 0.105 which is 66% of the overfishing threshold proxy ($F_{MSY\ proxy} = 0.1599$).

Note that if the missing CAA is left blank in this analysis there were slight differences compared to using a pooled Age Length key. Without the 2020 CAA, the retrospective pattern is better but doesn't use the length compositions currently in hand.

Based on this updated assessment, the white hake stock is not overfished and overfishing is not occurring.

As with other stocks, the Panel recommends that the pooled age length key be used rather than simply leaving the missing 2020 blank in the reference point calculations.

5. Conduct short-term stock projections when appropriate.

This **TOR** was satisfactorily addressed.

Short term projections of catch and **SSB** were derived by sampling from a cumulative distribution function of recruitment estimates from **ASAP** from 1995–2019. The annual fishery selectivity, maturity ogive, and mean weights at age used in the projection are the most recent 5-year (2017–2021) averages. The numbers-at-age used to start the projections were adjusted for retrospective bias using age-specific ρ estimates.

The estimated catch for 2022 is 1964 **mt**, which results in catch advice of 2347, 2258, and 2285 **mt** for 2023–2025, respectively.

*While the projection technique appears appropriate, the Panel recommends averaging **CAA/WAA** over the most recent five years of data without the inclusion of 2020.*

6. Respond to any review panel comments or **SSC** concerns from the most recent prior research or management track assessment.

This **TOR** was partially addressed, as the assessor presented their research recommendations but not those arising from previous assessment reviews or from the **SSC**.

This Panel has several research recommendations for potential improvement of the assessment:

- *The highest priority is to complete aging of collections from various sources with Fishery Observer collections being the most important (it provides missing information on catch). Then, aging of collections from the **ME/NH** and shrimp surveys should continue.*
- *Evaluate whether there is any difference between fishery and survey Weights-At-Age.*
- *Using a recruitment time series reaching back to 1995 may be inappropriate and **NMFS** should determine whether there is a more appropriate recruitment stanza for the stock. A change point analysis might be helpful.*
- *At the time of the next Management Track assessment, **NMFS** should reevaluate use of the shrimp survey data (is the improvement to the retrospective pattern an artifact of adding another index?)*
- *Reevaluate the utility of the Bottom Longline Survey in the assessment once additional years of data are available.*
- *Consider splitting the two **NEFSC** Bottom Trawl Surveys into separate **Albatross** and **Bigelow** indices.*
- *Evaluate the sensitivity of recruitment to the **CV** used (0.5).*


References:

NEFSC. 2013. 56th Northeast Regional Stock Assessment Workshop (SAW 56) Assessment Report. US Dept Commer, NOAA Fisheries, Northeast Fish Sci Cent Ref Doc. 13-10; 868p. [CRD13-10](#)

NEFSC. 2022. Operational Assessment of 14 Northeast Groundfish Stocks Updated Through 2018. US Dept Commer, NOAA Fisheries, Northeast Fish Sci Cent Ref Doc. 22-06; 227p. [CRD22-06](#)



Brian Gay of Millsboro, Delaware holding the record-setting white hake he caught in 2019. Photo credit: Maryland Department of Natural Resources.



Urophycis tenuis, White Hake.




Figure 29: Trends in spawning stock biomass of white hake between 1963 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $SSB_{Threshold}$ ($\frac{1}{2}SSB_{MSY proxy}$; horizontal dashed line) as well as SSB_{Target} ($SSB_{MSY proxy}$; horizontal dotted line) based on the 2022 assessment. Biomass was adjusted for a retrospective pattern and the adjustment is shown in red. The approximate 90% log-normal confidence intervals are shown.




Figure 30: Trends in the fully selected fishing mortality (F_{Full}) of white hake between 1963 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $F_{Threshold}$ ($F_{MSY proxy} = 0.1605$; horizontal dashed line). based on the 2022 assessment. The F_{Full} was adjusted for a retrospective pattern and the adjustment is shown in red. The approximate 90% log-normal confidence intervals are shown.




Figure 31: Trends in Recruits (age-1) (000s) of white hake between 1963 and 2021 from the current (solid line) and previous (dashed line) assessment. The approximate 90% log-normal confidence intervals are shown.




Figure 32: Total catch of white hake between 1963 and 2021 by fleet (commercial, recreational, or Canadian) and disposition (landings and discards).




Figure 33: Indices of biomass for white hake between 1963 and 2022 for the Northeast Fisheries Science Center (NEFSC) spring and fall bottom trawl surveys and the ASMFC shrimp survey. The approximate 90% log-normal confidence intervals are shown.

9. NORTHERN MONKFISH

Jon Deroba

*This assessment of the Northern Monkfish (*Lophius americanus*) stock is a Management Track assessment that updates the existing 2019 ‘Plan B’ assessment. This assessment report provides updated fishery catch data, the averaged NMFS spring and fall survey indices used in the ‘Plan B’ assessment, and the multiplier used to provide catch advice that results from the lsmooth approach used as the ‘Plan B’. Reference points were not applicable and stock status is unknown.*

State of Stock: Based on this Management Track assessment, Northern Monkfish (*Lophius americanus*) stock status is unknown.

Table 21: Catch table for Northern Monkfish. All weights are in (mt) .

	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Commercial discards	493	459	484	572	734	840	1,253	1,080	723	802
Commercial landings	4,081	3,355	3,434	4,086	4,723	7,105	6,009	6,084	5,587	5,121
Catch for Assessment	4,574	3,814	3,918	4,658	5,457	7,945	7,262	7,163	6,310	5,923

Table 22: References points are unavailable and stock status is unknown.

	2019	2022
$F_{MSY\ proxy}$	NA	NA
SSB_{MSY} (mt)	NA	NA
MSY (mt)	NA	NA
Overfishing	Unknown	Unknown
Overfished	Unknown	Unknown

Projections: Short term projections were not conducted due to lack of an analytical assessment. The catch multiplier resulting from the lsmooth approach equaled 0.829.

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

The largest source of uncertainty in this stock assessment is lack of a reliable aging method. Finding a reliable aging method seems unlikely. Monkfish migratory patterns also add uncertainty to the assessment and may complicate interpretation of indices of abundance.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted SSB or F_{Full} lies outside of the approximate joint confidence region for SSB and F_{Full}).

Not applicable.

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?

Not applicable.

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the effect these changes had on the assessment and stock status.

The assumed discard mortality rate of monkfish caught using scallop dredges was lowered from 100% to 64%. This change was based on a recent tagging study (Weissman et al., 2021).

- If the stock status has changed a lot since the previous assessment, explain why this occurred.

Not applicable.

- Provide qualitative statements describing the condition of the stock that relate to stock status.


*All indices of abundance reported in the stock assessment, including those not used in the *lsmooth* approach, showed declines of varying degrees over the past 3–5 years.*

- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.

The assessment could be most improved through development of an aging method, or exploration of a simpler two-stage, analytical assessment model.

- Are there other important issues?

The effects of climate change on monkfish biology and movement are likely to become increasingly important.



Lophius americanus, monkfish.

9.1. Reviewer Comments: Northern Monkfish

The 2022 assessment for the northern stock of monkfish (*Lophius piscatorius*) updates the 2019 assessment (NEFSC 2020)¹⁹ with additional commercial fishery catch data through 2021, and research survey indices of abundance and area-swept biomass through 2022.

An analytic assessment was not possible due to the lack of a reliable aging methodology. As a result, the ‘lsmooth’ (previously PlanBsmooth; Legault et al., in press)²⁰ approach used in the 2020 assessment was updated for this management track assessment. This ‘lsmooth’ approach re-scales the NMFS spring and fall BTS by their respective means (i.e., so each time series has mean equal to one) and averages the fall observation in year y with the spring observation in year $y + 1$ to create a single time series for analysis. A Loess-smooth is then applied to the combined time series, and a log-linear regression fit to the most recent three years of index predictions from the Loess fit. The slope of the regression provides a direction and rate of change in the indices that is multiplied by recent catch to provide catch advice. However, neither of the 2020 bottom trawl surveys were available. Consequently, the preferred approach was to use a combined spring and fall BTS time series with the missing 2020 observations replaced with the mean of the 2019 and 2021 observations. Using this method, the multiplier was 0.829 in the North.

An ‘lsmooth’ assessment does not allow for the estimation of reference points (i.e., F_{MSY} , and SSB_{MSY} cannot be determined). Therefore, the status of the stock relative to overfishing and being overfished must be unknown.

Short term projections are not possible using the ‘lsmooth’ approach.

The Panel spent considerable time discussing the appropriate term which the multiplier should be applied against: ABC or catch. The former has been the practice since the ‘lsmooth’ approach was first applied to monkfish and moving to catch would result in a major shift in catch advice. Applying the multiplier against the catch would result in a significant decrease in ABC advice. Estimates of area-swept minimum biomass developed from the chain sweep study indicate a high biomass from what is observed in the BTS but follow the same trends. On the other hand, the ‘lsmooth’ approach was designed to be applied to catch and is derived from catch data. Other index methods also are based on catch. Thus, application of the multiplier to catch is more consistent with lsmooth’s design and other index-based methods. Ultimately the group could not reach a consensus decision, though a majority supported the application of the multiplier against catch.

The Panel also considered whether stock status should be considered unknown. Given that the current stock status is based on a failed assessment, and that the ‘lsmooth’ approach does not generate reference points, the Panel strongly recommended listing stock status as unknown.

The Panel had several research recommendations:

- *Both the shrimp and scallop survey indices should be considered for inclusion in future assessments.*

¹⁹NEFSC 2020. Operational assessment of the black sea bass, scup, bluefish, and monkfish stocks, updated through 2018. NEFSC Ref Doc 20-01; 160p. [CRD20-01](#)

²⁰Legault, C.M., J. Wiedenmann, J.J. Deroba, G. Fay, T.J. Miller, E.N. Brooks, R.J. Bell, J.A. Langan, J.M. Courneane, A.W. Jones, and B. Muffley. 2022. Data Rich but Model Resistant: An Evaluation of data-limited methods to manage fisheries with failed age-based stock assessments. Canadian Journal of Fisheries and Aquatic Sciences. [CJFAS2022-0045](#); <https://github.com/cmlegault/PlanBsmooth>

- *Given the lack of success developing an aging technique, NMFS should not continue to pursue this avenue of research; consider estimating growth through cohort tracking.*
- *Given the lack of growth information on Monkfish, it was recommended the analyst explore a Simple Delay–Difference Model as a potential modeling approach relative to the ‘lsmooth’ method.*
- *Other Data Limited methods should also be considered for the assessment.*
- *A better understanding of stock structure (beyond North and South) could improve the assessment effort*
- *Reconsider the catchability coefficient of the chain swept estimates and how this applies to separate surveys.*

The Panel concluded that the 2022 assessment update for northern stock of monkfish fulfilled the recommendations of the AOP and is technically sufficient to provide scientific advice and meets the Terms of Reference for the stock’s assessment. It does not provide sufficient information to evaluate stock status. The assessment represents Best Scientific Information Available (BSIA) for this stock for management purposes.

References:

Weissman, A., Knotek, R., Mandelman, J., Rudders, D., Roman, S., and Sulikowski, J. 2021. Determining discard mortality of monkfish in a sea scallop dredge fishery. *North American Journal of Fisheries Management* 41: 856–870. [NAFM:10603](#)



Monkfish on the measuring table.




Figure 34: Total catch of Northern Monkfish by fleet (commercial, recreational, or Canadian) and disposition (landings and discards).




Figure 35: Indices of biomass for the Northern Monkfish between 1990 and 2022 for the Northeast Fisheries Science Center (NEFSC) spring and fall bottom trawl surveys. The approximate 90% log-normal confidence intervals are shown.

10. SOUTHERN MONKFISH

Jon Deroba

*This assessment of the Southern Monkfish (*Lophius americanus*) stock is a Management Track assessment that updates the existing 2019 ‘Plan B’ assessment. This assessment report provides updated fishery catch data, the averaged NMFS spring and fall survey indices used in the ‘Plan B’ assessment, and the multiplier used to provide catch advice that results from the ‘lsmooth’ approach used as the ‘Plan B’. Reference points were not applicable and stock status is unknown.*

State of Stock: Based on this Management Track assessment, Southern Monkfish (*Lophius americanus*) stock status is unknown.

Table 23: Catch table for Southern Monkfish. All weights are in (mt) .

	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Commercial discards	1,962	1,372	1,188	919	2,114	3,544	3,476	3,358	2,295	2,340
Commercial landings	5,674	5,207	5,099	4,550	4,331	3,796	4,388	4,373	2,593	2,005
Catch for Assessment	7,636	6,579	6,287	5,468	6,445	7,339	7,864	7,732	4,887	4,346

Table 24: Reference points are unavailable and stock status is unknown.

	2019	2022
$F_{MSY\ proxy}$	NA	NA
SSB_{MSY} (mt)	NA	NA
MSY (mt)	NA	NA
Overfishing	Unknown	Unknown
Overfished	Unknown	Unknown

Projections: Short term projections were not conducted due to lack of an analytical assessment. The catch multiplier resulting from the ‘lsmooth’ approach equaled 0.646.

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

The largest source of uncertainty in this stock assessment is lack of a reliable aging method. Finding a reliable aging method seems unlikely. Monkfish migratory patterns also add uncertainty to the assessment and may complicate interpretation of indices of abundance.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted SSB or F_{Full} lies outside of the approximate joint confidence region for SSB and F_{Full}).

Not applicable.

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?

Not applicable.

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the effect these changes had on the assessment and stock status.

The assumed discard mortality rate of monkfish caught using scallop dredges was lowered from 100% to 64%. This change was based on a recent tagging study (Weissman et al., 2021).

- If the stock status has changed a lot since the previous assessment, explain why this occurred.

Not applicable.

- Provide qualitative statements describing the condition of the stock that relate to stock status.


All indices of abundance reported in the stock assessment, including those not used in the 'smooth' approach, showed declines of varying degrees over the past 3–5 years.

- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.

The assessment could be most improved through development of an aging method, or exploration of a simpler two-stage, analytical assessment model.

- Are there other important issues?

The effects of climate change on monkfish biology and movement are likely to become increasingly important.



Lophius americanus, monkfish.

10.1. Reviewer Comments: Southern Monkfish

The 2022 assessment for the southern stock of monkfish (*Lophius piscatorius*) updates the 2019 assessment (NEFSC 2020)²¹ with additional commercial fishery catch data through 2021, and research survey indices of abundance and area-swept biomass through 2022.

An analytic assessment was not possible due to the lack of a reliable aging methodology. As a result, the ‘lsmooth’ (previously PlanBsmooth; Legault et al., in press)²² approach used in the 2020 assessment was updated for this management track assessment. This ‘lsmooth’ approach re-scales the NMFS spring and fall BTS by their respective means (i.e., so each time series has mean equal to one) and averages the fall observation in year y with the spring observation in year $y + 1$ to create a single time series for analysis. A Loess-smooth is then applied to the combined time series, and a log-linear regression fit to the most recent three years of index predictions from the Loess fit. The slope of the regression provides a direction and rate of change in the indices that is multiplied by recent catch to provide catch advice. However, neither of the 2020 bottom trawl surveys were available. Consequently, the preferred approach was to use a combined spring and fall BTS time series with the missing 2020 observations replaced with the mean of the 2019 and 2021 observations. Using this method, the multiplier was 0.646 in the south.

An ‘lsmooth’ assessment does not allow for the estimation of reference points (i.e., F_{MSY} , and SSB_{MSY} cannot be determined). Therefore, the status of the stock relative to overfishing and being overfished must be unknown.

Short term projections are not possible using the ‘lsmooth’ approach.

The Panel spent considerable time discussing the appropriate term which the multiplier should be applied against: ABC or catch. The former has been the practice since the ‘lsmooth’ approach was first applied to monkfish and moving to catch would result in a major shift in catch advice. Applying the multiplier against the catch would result in a significant decrease in ABC advice. Estimates of area-swept minimum biomass developed from the chain sweep study indicate a high biomass from what is observed in the BTS but follow the same trends. On the other hand, the ‘lsmooth’ approach was designed to be applied to catch and is derived from catch data. Other index methods also are based on catch, rather than ABC. Thus, application of the multiplier to catch is more consistent with lsmooth’s design and other index based methods. Ultimately the group could not reach a consensus decision, though a majority supported the application of the multiplier against catch.

The Panel also considered whether stock status should be considered unknown. Given that the current stock status is based on a failed assessment, and that the ‘lsmooth’ approach does not generate reference points, the Panel strongly recommended listing stock status as unknown.

The Panel had several research recommendations:

- *Both the shrimp and scallop survey indices should be considered for inclusion in future assessments.*

²¹NEFSC 2020. Operational assessment of the black sea bass, scup, bluefish, and monkfish stocks, updated through 2018. NEFSC Ref Doc 20-01; 160p. [CRD20-01](#)

²²Legault, C.M., J. Wiedenmann, J.J. Deroba, G. Fay, T.J. Miller, E.N. Brooks, R.J. Bell, J.A. Langan, J.M. Courneane, A.W. Jones, and B. Muffley. 2022. Data Rich but Model Resistant: An Evaluation of data-limited methods to manage fisheries with failed age-based stock assessments. Canadian Journal of Fisheries and Aquatic Sciences. [CJFAS2022-0045](#); <https://github.com/cmlegault/PlanBsmooth>

- *Given the lack of success developing an aging technique, NMFS should not continue to pursue this avenue of research further. Instead, NMFS should consider estimating growth through cohort tracking.*
- *Given the lack of growth information on Monkfish, it was recommended that the analyst explore a Simple Delay–Difference Model as one potential modeling approach in the next research track assessment.*
- *Other Data Limited methods should also be considered for that assessment.*
- *A better understanding of stock structure (beyond the border of Northern and Southern stocks) could improve the assessment effort.*
- *Reconsider the catchability coefficient of the chain swept estimates and how this applies to separate surveys.*

The Panel concluded that the 2022 assessment update for southern stock of monkfish fulfilled the recommendations of the AOP and is technically sufficient to provide scientific advice and meets the Terms of Reference for the stock’s assessment. It does not provide sufficient information to evaluate stock status. The assessment represents Best Scientific Information Available (BSIA) for this stock for management purposes.

References:

Weissman, A., Knotek, R., Mandelman, J., Rudders, D., Roman, S., and Sulikowski, J. 2021. Determining discard mortality of monkfish in a sea scallop dredge fishery. *North American Journal of Fisheries Management* 41: 856–870. [NAFM:10603](#)



Monkfish at gravelly sea bottom. Photo credit: iStock




Figure 36: Total catch of Southern Monkfish by fleet (commercial, recreational, or Canadian) and disposition (landings and discards).

NEFSC Spring and Fall Averaged




Figure 37: Indices of biomass for the Southern Monkfish between 1990 and 2022 for the Northeast Fisheries Science Center (NEFSC) spring and fall bottom trawl surveys. The approximate 90% log-normal confidence intervals are shown.

11. OCEAN POUT

Charles Adams

This assessment of the ocean pout (*Zoarces americanus*) stock is a level-1 management track assessment of the existing benchmark assessment (NEFSC 2008). Based on the previous 2020 management track assessment (NEFSC 2022) the stock was overfished, but overfishing was not occurring. This assessment updates commercial fishery catch data, research survey indices of abundance and exploitation ratios through 2021.

State of Stock: Based on this updated assessment, the ocean pout (*Zoarces americanus*) stock is overfished and overfishing is not occurring (Figures 38–39). Retrospective adjustments were not made to the model results. Biomass proxy (B) in 2021 was estimated to be 0.263 (kg/tow) which is 5% of the biomass target ($SSB_{MSY proxy} = 4.94$; Figure 38). The 2021 fully selected fishing mortality was estimated to be 0.234 which is 31% of the overfishing threshold proxy ($F_{MSY proxy} = 0.76$; Figure 39).

Table 25: Catch and status table for ocean pout. All weights are in (mt), survey biomass is in (kg/tow) and the exploitation ratio is catch/3-year moving average of NEFSC spring survey biomass index. Model results are from the current updated index assessment.

	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
	<i>Data</i>									
Commercial landings	0	0	0	0	0	0	0	0	0	0
Commercial discards	94	68	74	63	49	42	41	79	18	61
Other landings	0	0	0	0	0	0	0	0	0	0
Catch for Assessment	94	68	74	63	49	42	41	79	18	61
	<i>Model Results</i>									
NEFSC 3-yr average spring survey	0.317	0.37	0.301	0.319	0.223	0.232	0.182	0.175	0.285	0.263
Exploitation Ratio	0.296	0.185	0.248	0.197	0.222	0.183	0.224	0.455	0.062	0.234

Table 26: Comparison of reference points estimated in the 2020 management track and from the current assessment update. The median 3-year moving average of NEFSC spring survey biomass index and median exploitation ratio during 1977–1985 are used as B_{MSY} and F_{MSY} proxies, respectively.

	2020	2022
$F_{MSY proxy}$	0.76	0.76
SSB_{MSY} (kg/tow)	4.94	4.94
MSY (mt)	3,754	3,754
Overfishing	No	No
Overfished	Yes	Yes

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

An important source of uncertainty is the stock has not responded to low catch as expected.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted SSB or F_{Full} lies outside of the approximate joint confidence region for SSB and F_{Full}).

The exploitation ratio does not allow estimation of a retrospective pattern.

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?

Projections are not available for the exploitation ratio.

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the effect these changes had on the assessment and stock status.

*The time series of **Bigelow** indices was recalculated using station-specific swept areas.*

*Supplemental Figure 6 (see **SASINF**) was presented to the Assessment Oversight Panel (**AOP**) on May 23, 2022; the **AOP** agreed that the differences were minor.*

*The data source for commercial landings changed to the Catch Accounting and Monitoring System (**CAMS**) beginning in 2020. However, given the no possession limit, the **AOP** agreed that this is not an issue.*

- If the stock status has changed a lot since the previous assessment, explain why this occurred.

Stock status has not changed since the previous assessment.

- Provide qualitative statements describing the condition of the stock that relate to stock status.

*Discards comprise most of the catch since the no possession regulation was implemented in May 2010. The **NEFSC** survey indices remain at near-record low levels; there are few large fish in the population. The ocean pout stock remains in poor condition.*

- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.

The assessment could be improved with studies that explore why this stock is not rebuilding as expected.

- Are there other important issues?

*The 2020 spring **NEFSC** survey was treated as missing for this assessment. Thus, the moving average was calculated as the mean of the 2021 and 2022 survey indices. It is worth noting that a similar approach is taken at the start of the time series, where the moving average for 1968 is calculated as the mean of the 1968 and 1969 survey indices.*

The Assessment Oversight Panel (AOP) determined that the 2022 management track assessment for ocean pout would be a Level 1 direct delivery on May 23, 2022. Shortly afterward, while reviewing the report from the October 2020 NEFMC SSC meeting, this statement was noted “The SSC had a thorough discussion about ocean pout, and contemplated stating that the OFL was unknown but decided not to do this. The reference point used for this stock ($F_{MSY} = 0.76$) was set several assessment cycles back and, as such, probably needs to be reexamined to determine whether it is still applicable”.

The Population Dynamics Branch convened a brainstorming session on July 6, 2022 to address this issue. It was concluded that:

- Survey indices for ocean pout remain at, or near, record low levels, and that since the implementation of the no possession limit in May 2010, catch has consisted almost exclusively of discards.*
- In spite of the no possession limit, stock size has not increased, suggesting that this stock may have entered a depensatory state.*
- There is no clear alternative to the current reference point proxies, which are based upon survey biomass trends and the exploitation history.*
- The current reference points should remain in place until more justifiable alternatives can be developed.*
- More generally, removal of biological reference points (BRPs) is not desirable unless there is a clear justification on why the BRPs are no longer appropriate for stock status determination.*

11.1. Reviewer Comments: Ocean pout

Ocean pout was not peer reviewed in fall of 2022.

References:

Northeast Fisheries Science Center (NEFSC). 2008. Assessment of 19 northeast groundfish stocks through 2007: report of the 3rd Groundfish Assessment Review Meeting (GARM III), Northeast Fisheries Science Center, Woods Hole, Massachusetts, August 4–8, 2008. US Dept Commer, NOAA Fisheries, Northeast Fish Sci Cent Ref Doc. 08-15; 884p.+xvii. [CRD08-15](#)

Northeast Fisheries Science Center (NEFSC). 2022. Fall management track assessments 2020. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 22-08; 168p. [CRD22-08](#)




Figure 38: Trends in biomass (kg/tow) of ocean pout between 1968 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $SSB_{Threshold}$ ($\frac{1}{2}SSB_{MSY proxy}$; horizontal dashed line) as well as SSB_{Target} ($SSB_{MSY proxy}$; horizontal dotted line) based on the 2022 assessment.




Figure 39: Trends in the exploitation ratio of ocean pout between 1968 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $F_{Threshold}$ ($F_{MSY proxy} = 0.76$; horizontal dashed line) based on the 2022 assessment.




Figure 40: Total catch of ocean pout between 1968 and 2021 by fleet (US and other) and disposition (landings and discards). Note that a no possession limit was put in place in May 2010.




Figure 41: Indices of biomass (kg/tow) for ocean pout between 1968 and 2022 for the Northeast Fisheries Science Center (NEFSC) spring bottom trawl survey. The approximate 90% log-normal confidence intervals are shown.

12. AMERICAN PLAICE

Larry Alade

This assessment of the American plaice (*Hippoglossoides platessoides*) stock is a Level 2 Management Track assessment of the existing 2022 research track assessment (NEFSC 2022). Based on the previous assessment the stock was not overfished and overfishing is not occurring. This assessment updates commercial fishery catch data, research survey indices of abundance, and the analytical WHAM assessment model and reference points through 2021. Additionally, stock projections have been updated through 2025.

State of Stock: Based on this Management Track assessment, the American plaice (*Hippoglossoides platessoides*) stock is not overfished and overfishing is not occurring (Figures 42–43). Retrospective adjustment patterns were minor and retrospective adjustments for terminal year estimates were not needed. Spawning stock biomass (*SSB*) in 2021 was estimated to be 18,809mt, corresponding to 99% of the biomass target ($SSB_{MSY\ proxy} = 19,051$; Figure 42). The 2021 fully selected fishing mortality was estimated to be 0.045, corresponding to 11% of the overfishing threshold proxy ($F_{MSY\ proxy} = 0.414$; Figure 43).

Table 27: Catch and status table for American plaice. All weights are in (mt), recruitment is in (000s), and F_{Full} is the fishing mortality on fully selected ages (ages 6–9). Model results are from the current WHAM assessment updated through 2021.

	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
	<i>Data</i>									
Commercial landings	1,462	1,297	1,239	1,231	1,009	1,134	1,087	970	594	632
Commercial discards	302	168	92	86	108	102	109	78	68	69
Total catch	1,765	1,465	1,331	1,317	1,117	1,235	1,196	1,048	662	701
	<i>Model Results</i>									
Spawning Stock Biomass	26,478	23,844	30,474	35,146	43,867	46,479	43,085	29,540	22,124	18,809
F_{Full}	0.065	0.061	0.046	0.045	0.033	0.031	0.029	0.039	0.033	0.045
Recruits (age-1)	50,075	71,522	108,274	29,879	54,799	15,927	65,028	43,527	60,999	79,235

Projections: Short-term projections were conducted in the WHAM model, which propagates uncertainty in the recruitment and survival processes. For projection specifications, the NEFMC Groundfish Plan Development Team supplied an interim catch estimate for 2022. The annual fishery selectivity, maturity ogive, and mean weights-at-age used in the projections represent the most recent 5-year averages, following the decisions made at the 2022 research track. The stock is projected to remain above the overfished threshold throughout the short-term projection period.

Table 28: Comparison of revised reference points estimated from the previous 2022 research track assessment and from the current management track assessment. An $F_{40\%SPR}$ proxy was used for the overfishing threshold. The median and 95% probability intervals are reported.

	2022 research track	2022
$F_{MSY\ proxy}$	0.43	0.41
SSB_{MSY} (mt)	19,268	19,051 (15,073–24,080)
MSY (mt)	6492	6203 (4891–7867)
Median recruits (age-1) (000s)	51,270	51,987
Overfishing	No	No
Overfished	No	No

Table 29: Short term projections of total fishery catch and spawning stock biomass for American plaice based on a harvest scenario of fishing at $F_{MSY\ proxy}$ between 2023 and 2025. Catch in 2022 was assumed to be 653 (mt).

Year	Catch (mt)	SSB (mt)	F_{Full}
2022	653	21529 (14381–32231)	0.036
Year	Catch (mt)	SSB (mt)	F_{Full}
2023	7316	22536 (14241–35663)	0.414
2024	6670	20324 (11898–34718)	0.414
2025	6105	18298 (10040–33349)	0.414

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

The most important source of uncertainty in this assessment is in the commercial landings age samples available for 2020 and 2021. The initial application of the 2020 and 2021 commercial landed age samples resulted in a significant decrease in fish weights-at-age that could not be explained biologically (i.e., cohort shrinkage). Inspection of growth data (i.e., the age-length keys) showed evidence of skewness in the length-at age distribution, particularly in 2021. There is very little confidence that the age samples from the commercial landings for 2020 and 2021 are representative of the landed catch as it may relate to a sampling issue. For the purpose of this assessment, the 2019 age samples were used to characterize the age composition of the commercial landed catch data for both 2020 and 2021. Given the decadal growth pattern in American plaice, this alternative approach to characterizing the age composition for the last two years of landings time series for the assessment is reasonably justified.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted SSB or F_{Full} lies outside of the approximate joint confidence region for SSB and F_{Full}).

The 7-year Mohn's ρ , relative to *SSB*, was -0.023 in the 2022 research track assessment and 0.084 in 2021 for the current Management Track assessment. The 7-year Mohn's ρ , relative to *F*, was 0.031 in the 2022 research track assessment and -0.065 in 2021 for the current Management Track assessment. The retrospective pattern for this assessment was considered to be minor because the ρ -adjusted estimates of 2021 *SSB* ($SSB_{\rho} = 17354$) and 2021 *F* ($F_{\rho} = 0.048$) were within the approximate 95% confidence intervals around *SSB* (13,229–26,742) and *F* (0.029–0.071). Consequently, a retrospective adjustment of spawning stock biomass or fishing mortality in 2021 was not required.

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?

Population projections for American plaice are well determined. The stock is not in a rebuilding plan.

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the effect these changes had on the assessment and stock status.

The commercial landings changed to the Catch Accounting and Monitoring System (CAMS) beginning in 2020. Supplementary information was presented to the Assessment Oversight Panel (AOP) on August 3rd, 2022; the AOP concurred that there were no notable differences between the AA tables and CAMS. Additionally, the input data for the research track WHAM model were revised to address two issues. The first was to correct for the appropriate survey timing for spring and fall NEFSC survey indices. Second, the mapping of the weights-at-age matrices to the correct survey index was also revised to match the correct survey. Bridge model runs were conducted and the impacts of these changes on model results were inconsequential.

- If the stock status has changed a lot since the previous assessment, explain why this occurred.

The stock status of American plaice has not changed since the previous assessment (NEFSC 2022).

- Provide qualitative statements describing the condition of the stock that relate to stock status.

The NEFSC fall and spring survey indices continue to show large declines in abundance and weight. This partly explains the continued decline in model estimates of SSB since 2019.

- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.

The peer-review panel from the 2022 research track identified a number of research recommendations, the most important of which is to explore the availability of observer age samples for characterizing the age composition of commercial discards. The current approach uses survey age-length keys. Recent communication with the NEFSC age and growth lab indicated there are some archived observer age samples but the number of samples appears to be very low.

- Are there other important issues?

As indicated earlier, the commercial landings age samples are a source of uncertainty in this assessment. Future evaluation of these samples is warranted to determine how best to use this information in future assessments. Supplementary material is available on the Stock Assessment Supplementary Information website (SASINF).

12.1. Reviewer Comments: American plaice

The 2022 assessment for American plaice (*Hippoglossoides platessoides*) is the Management Track assessment resulting from the 2022 Research Track assessment (NEFSC 2022)²³. This WHAM assessment updates commercial fishery catch data, research survey indices of abundance, and reference points through 2021. Stock projections have been updated through 2025.

Retrospective patterns were minor and retrospective adjustments for terminal year estimates were not needed. Spawning stock biomass in 2021 was estimated to be 18,809 mt, corresponding to 99% of the biomass target ($SSB_{MSY\ proxy} = 19,051\text{mt}$). The 2021 fully selected fishing mortality was estimated to be 0.045, corresponding to 11% of the overfishing threshold proxy ($F_{MSY\ proxy} = 0.414$).

Based on this management track assessment, the stock was not overfished and overfishing was not occurring.

Short-term projections were conducted in the WHAM model, which propagates uncertainty in the recruitment and survival processes. The annual fishery selectivity, maturity ogive, and mean weights-at-age used in the projections represent the most recent 5-year averages, following the decisions made at the 2022 research track. The stock is projected to remain above the overfished threshold throughout the short-term projection period.

The estimated catch for 2022 is 653 mt, which results in catch advice of 7316, 6670, and 6105 mt for 2023–2025, respectively.

The Panel focused much of their discussion on the choice of WAA in both the base model and the projections. Both the 2020 and 2021 catch WAA showed steep declines from the 2019 data. These declines were sufficiently large to cast doubt on the data. Various explanations were provided for the observed decrease of WAA, none of which could be easily resolved. The analyst considered several approaches to the analysis including using the observed WAA, using no WAA for the two years, and substituting 2019 WAA for the 2020–2021 WAA. Ultimately, the best solution appeared to be the latter. After considerable discussion, the Panel recommended that the 2019 WAA be used for both the base WHAM model and the projections.

The Panel also recommended that the cause of this phenomena be thoroughly reviewed. If this is a result of inadequate or biased Port Sampling of landings, it is an example of the problems that inadequate sampling will introduce in future assessments for this and other stocks. Here it appears to make the strong 2013 cohort very difficult to track.

The Panel endorses several of the Research Track recommendations:


- *Exploration of archived observer otolith samples for characterizing age composition of discards*
- *Investigation of approaches to compare models with and without environmental covariates.*
- *Investigation of alternative estimates of natural mortality*

²³Northeast Fisheries Science Center (In Progress). Report of the 2022 American plaice Research Track Assessment working group. Available at: [SASINF](#). [AmPlaiceWG](#)

The Panel concluded that the 2022 assessment update for American plaice fulfilled the recommendations of the AOP, is technically sufficient to evaluate stock status and provide scientific advice and meets the Terms of Reference for the stock's assessment. The assessment represents Best Scientific Information Available (BSIA) for this stock for management purposes.

References:

Northeast Fisheries Science Center (In Progress). Report of the 2022 American plaice Research Track Assessment working group. Available at: [SASINF](#). [AmPlaiceWG](#)



Archival drawing: American Plaice.




Figure 42: Trends in spawning stock biomass of American plaice between 1980 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $SSB_{Threshold}$ ($\frac{1}{2}SSB_{MSY proxy}$; horizontal dashed line) as well as SSB_{Target} ($SSB_{MSY proxy}$; horizontal dotted line) based on the 2022 assessment. The approximate 95% log-normal confidence intervals are shown.




Figure 43: Trends in the fully selected fishing mortality (F_{Full}) of American plaice between 1980 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $F_{\text{Threshold}}$ ($F_{\text{MSY proxy}} = 0.414$; horizontal dashed line). The approximate 95% lognormal confidence intervals are shown.




Figure 44: Trends in Recruits (age-1) (000s) of American plaice between 1980 and 2021 from the current (solid line) and previous (dashed line) assessment. The approximate 95% log-normal confidence intervals are shown.




Figure 45: Total fishery catch of American plaice between 1980 and 2021 by disposition (landings and discards).




Figure 46: Indices of biomass for American plaice between 1980 and 2021 for the Northeast Fisheries Science Center (NEFSC) spring and fall bottom trawl surveys. The approximate 95% log-normal confidence intervals are shown.

13. POLLOCK

Brian Linton

This assessment of the pollock (*Pollachius virens*) stock is a Level 3 Management Track assessment, updating the 2019 Operational Assessment (NEFSC 2022). This assessment updates commercial and recreational fishery catch data, research survey indices of abundance, the ASAP analytical models, and biological reference points through 2021. Additionally, stock projections have been updated through 2025. In what follows, there are two population assessment models brought forward from the 2019 Operational Assessment: the base model (dome-shaped survey selectivity), which is used to provide management advice; and the flat sel sensitivity model (flat-topped survey selectivity), which is included for the sole purpose of demonstrating the sensitivity of assessment results to survey selectivity assumptions. The most recent benchmark assessment of the pollock stock was in 2010 as part of the 50th Stock Assessment Review Committee (SARC 50; NEFSC 2010), which includes a full description of the model formulations.

State of Stock: The pollock (*Pollachius virens*) stock is not overfished and overfishing is not occurring (Figures 47–48). No retrospective adjustments were made to the model results. Spawning stock biomass (*SSB*) in 2021 was estimated to be 175,573 mt under the base model and 85,109 mt under the flat sel sensitivity model which is 191% and 150% (respectively) of the biomass target, an *SSB*_{MSY proxy} of *SSB* at *F*_{40%SPR} (92,130 mt and 56,817 mt; Figure 47). The 2021 age 5 to 7 average fishing mortality (*F*) was estimated to be 0.052 under the base model and 0.092 under the flat sel sensitivity model, which is 22% and 39% (respectively) of the overfishing threshold, an *F*_{MSY proxy} of *F*_{40%SPR} (0.235 and 0.237; Figure 48).

Table 30: Catch and status table for pollock. All weights are in (mt), recruitment is in (000s), and *F*_{AVG} is the age 5 to 7 average *F*. Unadjusted *SSB* and *F* estimates are reported. Model results are from the current base model and flat sel sensitivity model.

	2014	2015	2016	2017	2018	2019	2020	2021
	<i>Data</i>							
Commercial landings	4,545	3,046	2,582	3,249	3,078	3,167	3,527	3,369
Commercial discards	135	155	97	49	70	154	174	99
Recreational landings	1,504	750	1,028	1,239	684	359	766	327
Recreational discards	2,429	2,186	1,519	2,059	940	584	765	728
Catch for Assessment	8,613	6,138	5,226	6,597	4,772	4,265	5,231	4,522
	<i>Model Results (base)</i>							
Spawning Stock Biomass	159022	180571	183767	200551	207766	177261	187710	175573
<i>F</i> _{AVG}	0.104	0.066	0.051	0.056	0.038	0.036	0.047	0.052
Recruits (age-1)	42577	29008	19866	19446	20370	18831	28368	12833
	<i>Model Results (flat sel sensitivity)</i>							
Spawning Stock Biomass	65427	75376	80568	89402	92958	83941	90307	85109
<i>F</i> _{AVG}	0.211	0.133	0.101	0.11	0.073	0.066	0.086	0.092
Recruits (age-1)	25325	17450	12108	12098	12924	12154	18413	8270

Table 31: Comparison of biological reference points for pollock estimated in the 2019 assessment and from the current base model and flat sel sensitivity model. An F_{MSY} proxy of $F_{40\%SPR}$ was used for the overfishing threshold, and was based on yield per recruit analysis. F_{MSY} is reported as the age 5 to 7 average F . Recruits represent the median of the predicted recruits. Intervals shown are 5th and 95th percentiles.

	2019 base	2019 flat sel sensitivity	base	flat sel sensitivity
F_{MSY}	0.272	0.260	0.235	0.237
SSB_{MSY} (mt)	124,639	70,721	92,130 (76,146–112,774)	56,817 (46,945–69,163)
MSY (mt)	19,856	12,007	12,519 (9,775–16,548)	8,238 (6,489–10,695)
Median recruits (age-1) (000s)	25,312	14,503	20,390	12,578
Overfishing	No	No	No	No
Overfished	No	No	No	No

Projections: Short term projections of median total fishery yield and spawning stock biomass for pollock were conducted based on a harvest scenario of fishing at an F_{MSY} proxy of $F_{40\%SPR}$ between 2023 and 2025. Catch in 2022 has been estimated at 3,959 (mt). Recruitments were sampled from a cumulative distribution function derived from ASAP estimated age-1 recruitment between 1970 and 2019. Recruitments in 2020 and 2021 were not included due to uncertainty in those estimates. The annual fishery selectivity, natural mortality, maturity ogive, and mean weights used in projections are the most recent 5-year averages. Retrospective adjusted SSB and age 5 to 7 average F in 2021 fell inside the 90% confidence intervals of the unadjusted 2021 value under the base model (Figures 47–48). Retrospective adjusted SSB and age 5 to 7 average F in 2021 fell inside the 90% confidence intervals of the unadjusted 2021 values under the flat sel sensitivity model (Figures 47–48). Therefore, no retrospective adjustments were made to the initial numbers-at-age in the projections for the base and flat sel sensitivity models.

Table 32: Unadjusted short term projections of median total fishery yield and spawning stock biomass for pollock from the current base model and flat sel sensitivity model based on a harvest scenario of fishing at an F_{MSY} proxy of $F_{40\%SPR}$ between 2023 and 2025. Catch in 2022 has been estimated at 3,959 (mt). F_{AVG} is the age 5 to 7 average F .

Year	Catch (mt)	SSB (mt) base	F_{AVG}	Catch (mt)	SSB (mt) flat sel sensitivity	F_{AVG}
2022	3,959	221,460	0.042	3,959	106,896	0.072
Year	Catch (mt)	SSB (mt) base	F_{AVG}	Catch (mt)	SSB (mt) flat sel sensitivity	F_{AVG}
2023	19,617	213,579	0.235	11,760	105,307	0.237
2024	17,546	189,776	0.235	10,769	94,974	0.237
2025	16,245	170,132	0.235	10,168	86,279	0.237

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

One of the greatest sources of uncertainty in the pollock assessment is selectivity, as the base model with dome-shaped survey and fishery selectivity curves implies the existence of a large cryptic biomass that neither current surveys nor the fishery can confirm. Assuming that survey selectivity is flat-topped leads to lower estimates of SSB and higher estimates of F . Stock status is insensitive to the shape of the survey selectivity patterns at older ages. Short term projection results differ based on the shape of the survey selectivity patterns at older ages. After the assessment, a consequence analysis is normally conducted to evaluate the management implications of the survey selectivity assumptions.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted SSB or F_{AVG} lies outside of the approximate joint confidence region for SSB and F_{AVG}).

The 7-year Mohn's ρ , relative to SSB , was 0.301 under the base model and 0.579 under the flat sel sensitivity model in the 2019 assessment and was 0.163 and 0.376, respectively, in 2021. The 7-year Mohn's ρ , relative now to F , was -0.282 under the base model and -0.389 under the flat sel sensitivity model in the 2019 assessment and was -0.154 and -0.263 , respectively, in 2021. There was a minor retrospective pattern for the base model because the ρ -adjusted estimates of 2021 SSB ($SSB_{\rho} = 175,573$ mt) and 2021 F ($F_{\rho} = 0.052$) were inside the approximate 90% confidence region around SSB (74,991–276,155 mt) and F (0.026–0.077). There was a minor retrospective pattern for the flat sel sensitivity model because the ρ -adjusted estimates of 2021 SSB ($SSB_{\rho} = 85,109$ mt) and 2021 F ($F_{\rho} = 0.092$) were inside the approximate 90% confidence region around SSB (46,729–123,490 mt) and F (0.047–0.137). No retrospective adjustment was made for the determination of stock status and for projections of catch in 2023 for both the base and flat sel sensitivity models.

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?

Population projections for pollock appear to be reasonably well determined for both the base and flat sel sensitivity models. The stock is not in a rebuilding plan.

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the affect these changes had on the assessment and stock status.

Three changes were explored in the current pollock assessment. First, tow-specific swept-area spring and fall bottom trawl survey indices were include in the assessment models. The tow-specific swept-area survey indices were similar to the traditional survey indices. The change in survey indices does not appear to have affected the assessment results. Second, based on a recommendation from the 2019 peer review panel, the separate commercial and recreational fleets were combined into a single fleet to reduce over-parameterization and improve model convergence. Combining the fishing fleets did improve model convergence and stability, while producing similar SSB and F estimates as the two-fleet model. Third, based on a recommendation from the 2019

peer review panel, the first year of the assessment was changed from 1970 to 1981, the first year of recreational catch data. The peer review panel was concerned that the sudden introduction of the new calibrated **MRIP** recreational catch estimates in 1981 might cause problems for the models. When the 1981 start year was applied to the one-fleet model, model stability and diagnostics were similar to the 1970-start-year model, but the stock was estimated to be less productive. The one-fleet 1970-start-year model was put forward as the base model, because dropping the first 11 years of data did not lead to a significant improvement in model diagnostics.

- If the stock status has changed a lot since the previous assessment, explain why this occurred.
Stock status based on the base and flat sel sensitivity models has not changed since the previous assessment.
- Provide qualitative statements describing the condition of the stock that relate to stock status.
Total removals of pollock have declined since 2013. The spring survey index increased from 2013 to 2018, decreased in 2019, and has remained relatively constant since that time. The fall survey index decreased from 2014 to 2017, and has remained relatively constant since that time. Fishery and survey data suggest the existence of a relatively strong 2013 year class, which has recruited to the commercial fishery. Survey data suggests that older fish have begun to reappear in the stock since the late-1990s.
- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.
The pollock assessment could be improved with additional studies on gear selectivity. These studies could cover topics such as physical selectivity (e.g., multi-mesh gillnet), behavior (e.g., swimming endurance, escape behavior), geographic and vertical distribution by size and age, tag-recovery at size and age, and evaluating information on length-specific selectivity at older ages. A satellite tagging project has been funded, which may provide information on the vertical distribution of pollock in the water column.
- Are there other important issues?
*The assessment plan for pollock that was presented at the **AOP** meeting included the development and evaluation of a historic recreational catch time series, 1970–1980. Construction of the historic catch series was not completed, due to time constraints. The historic recreational catch scenario could be explored in a future assessment.*



Pollachius virens, Atlantic Pollock.

13.1. Reviewer Comments: Pollock

The 2022 assessment of the pollock (*Pollachius virens*) stock updates the 2019 ASAP assessment (NEFSC 2022)²⁴ with additional commercial and recreational fishery catch data, research survey indices of abundance, and biological reference points through 2021. Stock projections have been updated through 2025.

The Panel concluded that the 2022 assessment update for pollock fulfilled the recommendations of the AOP, is technically sufficient to evaluate stock status and provide scientific advice and meets the Terms of Reference for the stock's assessment. The assessment represents Best Scientific Information Available (BSIA) for this stock for management purposes.

Terms of Reference: Pollock

1. Estimate catch from all sources including landings and discards.

This TOR was satisfactorily addressed.

Commercial and recreational discards, landings, and age composition were updated through 2021. Total removals in 2021 were estimated to be 4,522 mt which represents 24% of the sub-ACL for the stock (18,549 mt).

2. Evaluate indices used in the assessment (e.g., indices of relative or absolute abundance, recruitment, state surveys, age-length data, etc.).

This TOR was satisfactorily addressed.

Spring and fall tow-specific swept area bottom NEFSC trawl survey indices as well as survey maturities, and weights at age were updated through 2021 (2020 surveys were not conducted).

Panel recommended continued research into survey selectivity including the satellite tagging project and pollock swimming speed studies that could help support or refute the dome shaped selectivity. Consideration should also be given to splitting the Albatross IV and Bigelow survey indices.

3. Estimate annual fishing mortality, recruitment and stock biomass (both total and spawning stock) as possible (depending on the assessment method) for the time series using the approved assessment method and estimate their uncertainty. Include retrospective analyses if possible (both historical and within-model) to allow a comparison with previous assessment results and projections, and to examine model fit.

(a) Include bridge runs to sequentially document each change from the previously accepted model to the updated model proposed for this peer review.

(b) Prepare a backup assessment approach that would serve as an alternative for providing scientific advice to management if the analytical assessment were to not pass review.

²⁴NEFSC. 2022. Operational Assessment of 14 Northeast Groundfish Stocks Updated Through 2018. US Dept Commer, NOAA Fisheries, Northeast Fish Sci Cent Ref Doc. 22-06; 227p. [CRD22-06](#)

This TOR was satisfactorily addressed.

The most recent benchmark assessment of the pollock stock was in 2010 as part of the 50th Stock Assessment Review Committee (NEFSC 2010)²⁵, which includes a full description of the ASAP model formulations. Two population assessment models were brought forward from the 2019 Operational Assessment: the base model (dome-shaped survey selectivity), which is used to provide management advice; and the flat-topped survey selectivity model (flat sel), which is included for the sole purpose of demonstrating the sensitivity of assessment results to survey selectivity assumptions. A bridge run comparing the 2019 and 2022 base models (2-Fleet Start year 1970) with the latter updated to the 2021 terminal year was run and showed that the 2013 year class was smaller than estimated in the 2019 assessment (and reduced terminal year SSB with little effect on F).

The analyst further considered four permutations of these models: 1-Fleet versus 2-Fleet and Start Year 1970 versus 1981. The 1-Fleet 1970-Start-Year model was selected as the 2022 base model, because the 1-Fleet model improved model convergence, and there was no significant improvement in model fit or performance to justify dropping the first 11 years of data from the assessment. A base model was obtained by fine-tuning the 1-Fleet 1970-Start-Year model to improve RMSE for each set of residuals.

The base model assumes dome-shaped fishery and survey selectivities, and this results in a cryptic biomass of older fish that cannot be confirmed by the fishery or surveys. A sensitivity model with flat-top survey selectivities was carried forward through reference point calculations and short-term projections to evaluate model sensitivity to selectivity assumptions (Flat-top selectivity = selectivity for ages 6+ fixed at 1.0). Note that the flat sel sensitivity model is only to be used as a sensitivity analysis and is not to be used for catch advice.

The Panel recommended that the one fleet model beginning in 1970 with dome shaped selectivity was the better model. It has the lowest AIC and retrospective pattern; the 1970 start date provides more historic data; and the model tuning further improved the retrospective pattern.

Concern was, however, expressed about selecting selectivity time-blocks based on visual inspection of residuals. Without some external forcing mechanism (e.g., a management measure or fishery shift) this introduces significant uncertainty into the selectivity function.

Also, the tuning process which relaxed the CV on the surveys and catch while tightening the CV on discards seemed counterintuitive, given discards are generally less well known than the catch or survey data.

The model results of SSB are very sensitive to the starting year (1970~1981) and to the selectivity assumptions (domed~flat), which indicate the scale of the model estimated stock size can be problematic. The % of fish of age 9+ in the CAA and survey age compositions are high based on the bubble plots. The estimated SSBs of age 9+ are higher than 40–60% of the stock's total SSB over the years even when flat selectivity was used. Future stock assessment may consider increasing the age of the plus group to see whether it can solve the scaling difficulty.

²⁵NEFSC. 2010. 50th Northeast Regional Stock Assessment Workshop (SAW 50) Assessment Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 10-17; 844p. [CRD10-17](#)

4. Re-estimate or update the BRPs as defined by the management track level and recommend stock status. Also, provide qualitative descriptions of stock status based on simple indicators/metrics (e.g., age- and size-structure, temporal trends in population size or recruitment indices, etc.).

This TOR was satisfactorily addressed.

No retrospective adjustments were made to the model results. Spawning stock biomass (SSB) in 2021 was estimated to be 175,573 mt under the base model and 85,109 mt under the flat sel sensitivity model which is 191 and 150% (respectively) of the biomass target, an SSB_{MSY} proxy of SSB at $F_{40\%SPR}$ (92,130 and 56,817 mt). The 2021 age 5 to 7 average fishing mortality (F) was estimated to be 0.052 under the base model and 0.092 under the flat sel sensitivity model, which is 22% and 39% (respectively) of the overfishing threshold, an F_{MSY} proxy of $F_{40\%SPR}$ (0.235 and 0.237).

The stock is not overfished and overfishing is not occurring.

5. Conduct short-term stock projections when appropriate.

This TOR was satisfactorily addressed.

Short term projections of median total fishery yield and spawning stock biomass for pollock were conducted using agepro and based on a harvest scenario of fishing at an F_{MSY} proxy of $F_{40\%SPR}$ between 2023 and 2025. Recruitments were sampled from a cumulative distribution function derived from ASAP estimated age-1 recruitment between 1970 and 2019. Recruitments in 2020 and 2021 were not included due to uncertainty in those estimates. The annual fishery selectivity, natural mortality, maturity ogive, and mean weights used in projections are the most recent 5-year averages. No retrospective adjustments were made.

The estimated catch for 2022 is 3959 mt, which results in catch advice of 19614, 17546, and 16249 mt for 2023–2025, respectively.

6. Respond to any review panel comments or SSC concerns from the most recent prior research or management track assessment.

This TOR was satisfactorily addressed.

The analyst has successfully responded to four significant sets of recommendations from the 2019 review.

This Panel recommended research into:

- *Survey selectivity including the satellite tagging and pollock swimming speed studies that could help support or refute the dome shaped selectivity*
- *Splitting the F/V Albatross and F/V Bigelow survey indices.*
- *Increase the age of the plus age group.*

References:

Northeast Fisheries Science Center. 2010. 50th Northeast Regional Stock Assessment Workshop ([SAW 50](#)) Assessment Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 10-17; 844p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026. [CRD10-17](#).

Northeast Fisheries Science Center. 2022. Stock Assessment Update of 14 Northeast Groundfish Stocks Through 2018. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 22-06; 232p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026. [CRD22-06](#).



Pollock in a basket.




Figure 47: Estimated trends in the spawning stock biomass of pollock between 1970 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $SSB_{Threshold}$ ($\frac{1}{2}SSB_{MSY proxy}$; horizontal dashed line) as well as SSB_{Target} ($SSB_{MSY proxy}$; horizontal dotted line) based on the 2022 assessment models base (A) and flat sel sensitivity (B). The retrospective adjusted biomass is shown in red. The approximate 90% log-normal confidence intervals are shown.





Figure 48: Estimated trends in age 5 to 7 average F (F_{AVG}) of pollock between 1970 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $F_{Threshold}$ ($F_{MSY\ proxy}$; dashed line) based on the 2022 assessment models base (A) and flat sel sensitivity (B). The retrospective adjusted F_{AVG} is shown in red. The approximate 90% log-normal confidence intervals are shown.




Figure 49: Estimated trends in age-1 recruitment (000s) of pollock between 1970 and 2021 from the current (solid line) and previous (dashed line) assessment for the assessment models base (A) and flat sel sensitivity (B). The approximate 90% log-normal confidence intervals are shown.




Figure 50: Total catch of pollock between 1970 and 2021 by fleet (commercial, Canadian, distant water fleet, and recreational) and disposition (landings and discards).




Figure 51: Indices of abundance for pollock from the Northeast Fisheries Science Center (NEFSC) spring (1970 to 2022) and fall (1970 to 2021) bottom trawl surveys. The approximate 90% log-normal confidence intervals are shown.

14. WITCH FLOUNDER

Susan Wigley

This assessment of the witch flounder (*Glyptocephalus cynoglossus*) stock is a management track assessment of the existing 2019 assessment (NEFSC 2022). Based on the 2019 assessment the stock status was overfished and overfishing unknown, and stock condition was poor. This assessment updates commercial fishery catch data through 2021 (Table 33, Figure 54), and updates research survey biomass indices and the empirical approach assessment through 2021 (Figure 55). No stock projections can be computed using the empirical approach.

State of Stock: Based on this updated assessment, witch flounder (*Glyptocephalus cynoglossus*) recommended stock status cannot be determined analytically due to a lack of biological reference points associated with the empirical approach; stock condition remains poor. Retrospective adjustments were not made to the model results. The exploitable biomass in 2021 (defined as the arithmetic average of the 2021 NEFSC spring and 2020 NEFSC fall surveys population biomass estimates and converted to exploitable biomass (multiplied by 0.9) based on examination of survey and fishery selectivity patterns) was estimated to be 22,419 (mt) (Figure 52). The 2021 exploitation rate (2021 catch divided by 2021 exploitable biomass) was estimated to be 0.039 (Figure 53).

Table 33: Catch and model results table for witch flounder. All weights are in (mt). The exploitable biomass in year y is the arithmetic average of the year y NEFSC spring and year $y - 1$ NEFSC fall surveys then converted to exploitable biomass using 0.9. The exploitation rate is the year y catch divided by the year y exploitable biomass. Model results are from the current updated empirical approach assessment.

	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
	<i>Data</i>										
Commercial Landings	870	1,038	686	570	492	397	446	606	800	869	823
Commercial Discards	201	231	124	107	94	115	106	115	95	90	56
Catch for Assessment	1,072	1,270	811	676	586	512	552	722	894	959	879
	<i>Model Results</i>										
Exploitable Biomass	16,313	18,404	12,533	14,628	17,289	15,847	22,247	32,546	25,736	27,958	22,419
Exploitation Rate	0.066	0.069	0.065	0.046	0.034	0.032	0.025	0.022	0.035	0.034	0.039

Table 34: Comparison of reference points estimated in an earlier assessment and from the current assessment update.

	2019	2022
$F_{MSY proxy}$	NA	NA
SSB_{MSY} (mt)	NA	NA
MSY (mt)	NA	NA
Overfishing	Unknown	Unknown
Overfished	Yes	Yes

Projections: Short term projections cannot be computed using the empirical approach. The estimated 2022 exploitable biomass is 19,393 mt. Using the January 2017 NEFMC PDT SSC approach for catch advice, application of the mean exploitation rate of 5.4% (based on nine years, 2007–2015) to the 3 year (2020–2022) moving average of exploitable biomass (23,257mt) results in an estimated catch for 2023 of 1,256 mt.

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

Uncertainty in the catch has increased due to criminal convictions in a case involving catch misreporting.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted SSB or F_{Full} lies outside of the approximate joint confidence region for SSB and F_{Full}).

The model used to estimate status of this stock does not allow estimation of a retrospective pattern.

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?

Population projections for witch flounder are not computed. Catch advice is derived from applying a mean exploitation rate of 0.054 (based on nine years, 2007–2015) to the 3-year average (2020–2022) of the exploitable biomass. The change in mean exploitation rate from 0.049 (2019 assessment) to 0.054 emanated from the use of tow-specific swept area indices. The stock is in a revised rebuilding plan, rebuilding by 2043. Since 2019, the average survey biomass has declined despite catches being below the catch advice from the 2019 assessment.

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the effect these changes had on the assessment and stock status.

Recent landings and discards were updated and the time series of survey indices were updated using tow-specific swept area; however, this has no impact on the stock status. In the 2022 assessment of witch flounder, the catch efficiency analyses were directly incorporated into the assessment model. Estimates of population biomass used revised seasonal catchability coefficients that varied by year; the revised seasonal catchability coefficients had a minor impact on catch advice for 2023. The 2018 NEFSC fall survey stratum 30 was not sampled; survey indices were not adjusted because this stratum represents less than 1% of total expanded catch weight and has negligible impact on survey indices and swept area biomass. The 2020 NEFSC spring and fall survey values are missing because of Covid. The missing survey values were not replaced with an average because the method used for catch advice is an average of the 3 most recent years of exploitable biomass, so filling the missing surveys was not deemed necessary. The 3 year (2020–2022) moving average exploitable biomass was derived using 2020 (Fall 2019 survey), 2021 (Spring 2021 survey), and 2022 (the average of Fall 2021 and Spring 2022).

- If the stock status has changed a lot since the previous assessment, explain why this occurred.
No change in stock status has occurred for witch flounder since the previous assessment. Biological references points remain unknown.
- Provide qualitative statements describing the condition of the stock that relate to stock status.
The witch flounder stock condition remains poor. Fishery landings and survey catch by age indicate continued truncation of age structure and a reduction in the number of older fish in the population. NEFSC relative indices of abundance and biomass remain below their time series average. The decline in exploitable biomass from 2020 to 2022 is notable.
- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.
The witch flounder assessment could be improved with accurate catch statistics; catch statistics have been undermined by misreporting, as partially documented in the criminal case. Although not directly impacting the empirical approach assessment, the low commercial landings sampling in recent years impacts the ability to estimate numbers of fish landed at age and to track cohorts through time. It would be desirable to review the port and dockside monitoring sampling to better align sampling activities in accordance with market category landings and life history characteristics of witch flounder. Additional research recommendations are given in NEFSC 2017b.
- Are there other important issues?
The empirical approach does not incorporate age structure information. Consideration of incoming recruitment is critical for catch advice that supports stock rebuilding. Based on the surveys, there is no sign of a strong incoming year class.
This assessment and the 2019 assessment used revised seasonal catchability coefficients (q vary by year) in the estimates of population biomass. The 2016 and 2017 assessments applied a constant catchability coefficient (0.291).
Minimum estimates of scientific research removals of witch flounder ranged between 0.1 and 15.9mt, with an average of 1mt between 1963 and 2018. The NEFSC bottom trawl surveys, Massachusetts Division of Marine Fisheries inshore surveys, Atlantic States Marine Fisheries Commission summer shrimp surveys, and various Cooperative Research surveys (e.g., such as Industry-based surveys for cod and for yellowtail flounder) and gear studies have contributed to scientific research removals. The August 2016 Gear Efficiency Study removed 14.0mt of witch flounder. These removals would be included when an age-based assessment is conducted.

14.1. Reviewer Comments: Witch flounder

Witch flounder was not peer reviewed in fall of 2022.

References:

Northeast Fisheries Science Center. 2022. Stock Assessment Update of 14 Northeast Groundfish Stocks Through 2018. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 22-06; 227p. [CRD22-06](#)

Northeast Fisheries Science Center. 2017a. Operational Assessment of 19 Northeast Groundfish Stocks, Updated Through 2016. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 17-17; 259p.

Northeast Fisheries Science Center. 2017b. 62nd Northeast Regional Stock Assessment Workshop Assessment Report, Northeast Fisheries Science Center, Woods Hole, Massachusetts, January 2017. US Dept Commer, [NOAA](#) Fisheries, Northeast Fish Sci Cent Ref Doc. 17-03; 822p.

Northeast Fisheries Science Center. 2015. Operational Assessment of 20 Northeast Groundfish Stocks, Updated through 2014. US Dept Commer, [NOAA](#) Fisheries, Northeast Fish Sci Cent Ref Doc. 15-24; 251p. [CRD15-24](#)




Figure 52: Trends in exploitable biomass (mt) of witch flounder between 1968 and 2022 from the current assessment.




Figure 53: Trends in the exploitation rate (catch/exploitable biomass) of witch flounder between 1982 and 2021 from the current assessment.




Figure 54: Total catch of witch flounder between 1982 and 2021 by fleet (commercial) and disposition (landings or discards).




Figure 55: Indices of biomass for witch flounder between 1963 and 2022 for the Northeast Fisheries Science Center (NEFSC) spring and fall bottom trawl surveys. The approximate 90% log-normal confidence intervals are shown.

15. CAPE COD–GULF OF MAINE YELLOWTAIL FLOUNDER

Larry Alade

*This assessment of the Cape Cod–Gulf of Maine yellowtail flounder (*Limanda ferruginea*) stock is an operational assessment of the existing 2019 VPA assessment (Alade 2019). The last benchmark for this stock was in 2008 (Legault et al., 2008). Based on the previous assessment the stock was not overfished, and overfishing was not occurring. This 2022 assessment updates commercial fishery catch data, research survey indices of abundance, weights at age, and the analytical VPA assessment model and reference points through 2021. Additionally, stock projections have been updated through 2025.*

State of Stock: Based on this updated assessment, Cape Cod–Gulf of Maine yellowtail flounder (*Limanda ferruginea*) stock is not overfished and overfishing is not occurring (Figures 56–57). Retrospective adjustments were made to the model results. Spawning stock biomass (*SSB*) in 2021 was estimated to be 3,058 (mt) which is 100% of the biomass target ($SSB_{MSY\ proxy} = 3,068$; Figure 56). The 2021 fully selected fishing mortality was estimated to be 0.1035 which is 32% of the overfishing threshold proxy ($F_{MSY\ proxy} = 0.3204$; Figure 57).

Table 35: Catch and model results for Cape Cod–Gulf of Maine yellowtail flounder. All weights are in (mt), recruitment is in (000s) and F_{Full} is the average fishing mortality on ages (ages 4 and 5). Model results below are from the current updated VPA assessment without any retrospective adjustment.

	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
<i>Data</i>										
Commercial discards	146	86	54	45	66	50	45	44	35	71
Commercial landings	946	590	421	306	302	314	226	184	156	294
Total Catch for Assessment	1,092	676	475	351	368	365	271	228	192	365
<i>Model Results</i>										
Spawning Stock Biomass	1,039	725	705	964	1,126	1,216	1,299	2,119	3,873	5,987
F_{Full}	1.127	1.209	0.644	0.36	0.272	0.28	0.233	0.155	0.06	0.05
Recruits (age-1)	2,271	3,412	2,923	2,357	3,953	6,517	13,357	20,854	9,269	10,413

Projections: Short term projections of biomass were derived by sampling an empirical cumulative distribution function of 35 recruitment estimates from the VPA model results. The most recent two years (2021 and 2022) were not included in the series of recruitment values due to high uncertainty in these estimates. The annual fishery selectivity, maturity ogive, and mean weights at age used in projection are the most recent 5-year averages. Retrospective adjustments were applied in the projections.

Table 36: Comparison of reference points estimated in the previous assessment and from the current assessment update. An $F_{40\%SPR}$ proxy was used for the overfishing threshold and $SSB_{MSY proxy}$ was based on long-term stochastic projections.

	2019	2022
$F_{MSY proxy}$	0.3204	0.3201
SSB_{MSY} (mt)	3,439	3,068 (2,108–4,751)
MSY (mt)	1,138	1,008 (696–1,554)
Median recruits (age-1) (000s)	5,781	6,417
Overfishing	No	No
Overfished	No	No

Table 37: Short term projections of total fishery catch and spawning stock biomass for Cape Cod–Gulf of Maine yellowtail flounder based on a harvest scenario of fishing at $F_{MSY proxy}$ between 2024 and 2025. Catch in 2022 was assumed to be 350 (mt).

Year	Catch (mt)	SSB (mt)	F_{Full}
2022	350	4,334 (3,512–5,360)	0.077

Year	Catch (mt)	SSB (mt)	F_{Full}
2023	1,436 (1,129–1,784)	4,433 (3,475–5,534)	0.320
2024	1,197 (955–1,494)	3,666 (2,929–4,542)	0.320
2025	1,059 (828–1,434)	3,269 (2,546–4,469)	0.320

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

Retrospective patterns remain a source of uncertainty in the assessment. This has persisted for a number of years causing a decrease in estimates of spawning stock biomass (SSB) and an increase in fishing mortality (F) when more years of data are added. The magnitude of these retrospective biases in this assessment have notably increased for both F and SSB compared to the previous 2019 Management Track assessment. With the exception of the 2020 ME/NH fall survey index, another potential source of uncertainty is the missing 2020 fall (NEFSC, MA DMF) and spring (NEFSC, MA DMF and ME/NH) survey data indices in the model.

In this assessment, the 2020 survey indices were assigned as missing in the VPA model due to unavailable surveys data in 2020 as a result of Covid. The treatment of missing data in the VPA model was not based on any form of imputations but rather the model was allowed to generate a survey prediction based on neighboring observed values. However, the model fit to the missing 2020 survey indices does not contribute to the overall objection function in the model.

The VPA model assumes catch is known without error, which is the case for this assessment and certainly not a valid assumption. The VPA model framework provides very little opportunity to leverage data uncertainty into estimates of population quantities produced by the model.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted SSB or F_{Full} lies outside of the approximate joint confidence region for SSB and F_{Full}).

The 7-year Mohn's ρ , relative to SSB , was 0.30 in the 2019 assessment and was 0.96 in 2021. The 7-year Mohn's ρ , relative to F , was -0.15 in the 2019 assessment and was -0.52 in 2021. There was a major retrospective pattern for this assessment because the ρ -adjusted estimates of 2021 SSB ($SSB_{\rho} = 3058$) and 2021 F ($F_{\rho} = 0.1035$) were outside the approximate 90% confidence region around SSB (4,976–7,428) and F (0.04–0.06). A retrospective adjustment was made for both the determination of stock status and for projections of catch in 2023. The retrospective adjustment changed the 2021 SSB from 5,987 to 3,058 and the 2021 F_{Full} from 0.05 to 0.1035.

- Based on this stock assessment, are population projections well determined or uncertain?
Population projections for Cape Cod-Gulf of Maine yellowtail flounder are uncertain for reasons associated with the retrospective bias in this updated assessment. The 2021 estimates of SSB from this assessment is not within the bound of values projected in the 2019 Management Track assessment. The 2019 estimate of SSB from the current assessment is below the the 2019 ρ -adjusted SSB from the 2019 Management Track assessment, indicating the ρ -adjustment applied in 2019 assessment was not large enough.

- Describe any changes that were made to the current stock assessment (e.g., catch efficiency studies), beyond incorporating additional years of data and the effect these changes had on the assessment and stock status.

Minor changes in addition to the incorporation of new data were made to the Cape Cod–Gulf of Maine yellowtail flounder assessment for this update. The NEFSC spring and fall indices were revised from 2009 to 2022 to account for tow-specific area swept. The data source for commercial landings changed to the Catch Accounting and Monitoring System (CAMS) beginning in 2020 and were used to produce commercial landings estimates for 2020 and 2021.

In the previous 2019 Management track assessment (NEFSC, 2022), The 2019 spring MA DMF survey age composition was derived by borrowing from the 2019 spring NEFSC ALK due to unavailable 2019 spring MA DMF ages at the time of the assessment. In this assessment, the 2019 spring survey age composition was revised to use the MA DMF Age-length keys (ALK), consistent with the previous assessment and the benchmark formulation in 2008 (GARM III). A sensitivity run was conducted to evaluate the impact of this change. The revision to the 2019 spring MA DMF survey age composition resulted in little to no effect on the assessment results.

The incorporation of new data (2019–2021) to the model resulted in rescaling of SSB and F which partially explains the cause of retrospective pattern in the model.

- If the stock status has changed a lot since the previous assessment, explain why this occurred.
The stock status for Cape Cod–Gulf of Maine yellowtail flounder is now rebuilt due increases in the survey biomass. Based on this assessment, estimated SSB in 2021 is above both the $SSB_{Threshold}$ and SSB_{Target} . The stock is in a rebuilding plan with a rebuild date of 2023. Based on the the 2022 assessment, the stock is rebuilt and continues to be in the 3-year projections.
- Provide qualitative statements describing the condition of the stock that relate to stock status.
All indices has shown increases in recent years. $CCGM$ yellowtail stock show no truncation in the age structure. There has been some moderate expansion in the older age groups which is also supported by the surveys. There is an above average estimated 2018 and 2020 incoming year classes which has contributed to the increase in total biomass. Estimates of commercial catch continue are still levels compared to historical catches and consistent with increase in stock biomass.
- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.
The Cape Cod–Gulf of Maine yellowtail flounder assessment could be improved with a change in model platform that incorporates statistical fits and accounts for measures of uncertainty in the model. Additionally, this assessment could benefit from updated growth and maturity studies. The current maturity and growth parameters are based on $GARM III$ estimates ($NEFSC$ 2008) which are over a decade old. It should be noted that the Cape Cod-Gulf of Maine yellowtail assessment is currently undergoing a Research Track assessment, scheduled for 2024.
- Are there other important issues?
None.

15.1. Reviewer Comments: Cape Cod–Gulf of Maine yellowtail flounder

Cape Cod–Gulf of Maine yellowtail flounder was not peer reviewed in fall of 2022.

References:

Miller T.J., 2013. A comparison of hierarchical models for relative catch efficiency based on paired-gear data for U.S. northwest Atlantic fish stocks. *Canadian Journal of Fisheries and Aquatic Sciences* 70(9): 1306–1316. [2013-0136](#)

Miller T.J., Martin M., Politis P., Legault C.M., Blaylock J. 2017a. Some statistical approaches to combine paired observations of chain sweep and rockhopper gear and catches from **NEFSC** and **DFO** trawl surveys in estimating Georges Bank yellowtail flounder biomass. **TRAC** Working Paper 2017/XX; 36p.

Miller T.J., Richardson D.E., Politis P., Blaylock J. 2017b. **NEFSC** bottom trawl catch efficiency and biomass estimates for 2009–2017 for 8 flatfish stocks included in the 2017 Northeast Groundfish Operational Assessments. Working paper. National Marine Fisheries Service, Northeast Fisheries Science Center, Woods Hole, MA. September 11–15, 2017.

Miller T.J., Politis P., Blaylock J., Richardson D., Manderson J., Roebuck C. 2018. Relative efficiency of a chain sweep and the rockhopper sweep used for the **NEFSC** bottom trawl survey and chainsweep-based swept area biomass estimates for 11 flatfish stocks. **SAW 66** summer flounder Data/Model/Biological Reference Point (**BRP**) meeting. National Marine Fisheries Service, Northeast Fisheries Science Center, Woods Hole, MA. September 17–21, 2018.

Legault C., L. Alade, S. Cadrin, J. King, and S. Sherman. 2008. In Northeast Fisheries Science Center. 2008. Assessment of 19 Northeast Groundfish Stocks through 2007: Report of the 3rd Groundfish Assessment Review Meeting (**GARM III**), Northeast Fisheries Science Center, Woods Hole, Massachusetts, August 4–8, 2008. US Dept Commer, **NOAA** Fisheries, Northeast Fish Sci Cent Ref Doc. 08-15; 884p.+xvii. [CRD08-15](#)

Legault C., L. Alade, S. Emery, J. King, and S. Sherman. 2012. In Northeast Fisheries Science Center. 2012. Assessment or Data Updates of 13 Northeast Groundfish Stocks through 2010. US Dept Commer, **NOAA** Fisheries, Northeast Fish Sci Cent Ref Doc. 12-06.; 789p. [CRD12-06](#)

Alade, L. 2015. In Northeast Fisheries Science Center. 2015. Operational Assessment of 20 Northeast Groundfish Stocks, Updated Through 2014. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 15-24; 251p. [CRD15-24](#)

Alade, L. 2017. In Northeast Fisheries Science Center. 2017. Operational Assessment of 19 Northeast Groundfish Stocks, Updated Through 2016. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 15-17; 259p. [CRD15-17](#)

Northeast Fisheries Science Center, 2022. Stock Assessment Update of 14 Northeast Groundfish Stocks Through 2018. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 22-06; 232p. [CRD22-06](#)




Figure 56: Trends in spawning stock biomass of Cape Cod–Gulf of Maine yellowtail flounder between 1985 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $SSB_{Threshold}$ ($\frac{1}{2}SSB_{MSY proxy}$; horizontal dashed line) as well as SSB_{Target} ($SSB_{MSY proxy}$; horizontal dotted line) based on the 2022 assessment. Biomass was adjusted for a retrospective pattern and the adjustment is shown in red. The 90% bootstrap probability intervals are shown.




Figure 57: Trends in the fully selected fishing mortality (F_{Full}) of Cape Cod–Gulf of Maine yellowtail flounder between 1985 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $F_{\text{Threshold}}$ ($F_{\text{MSY proxy}} = 0.3204$; horizontal dashed line). F_{Full} was adjusted for a retrospective pattern and the adjustment is shown in red based on the 2022 assessment. The 90% bootstrap probability intervals are shown.




Figure 58: Trends in Recruits (age-1) (000s) of Cape Cod-Gulf of Maine yellowtail flounder between 1985 and 2021 from the current (solid line) and previous (dashed line) assessment. The 90% bootstrap probability intervals are shown.




Figure 59: Total catch of Cape Cod–Gulf of Maine yellowtail flounder between 1985 and 2021 by disposition (landings and discards).




Figure 60: Indices of biomass for the Cape Cod–Gulf of Maine yellowtail flounder between 1985 and 2022 for the Northeast Fisheries Science Center (NEFSC) spring and fall bottom trawl surveys, Massachusetts Department of Marine Fisheries (MADMF) inshore state spring and fall bottom trawl surveys, and the Maine–New Hampshire inshore state spring and fall state surveys. The 90% bootstrap probability intervals are shown.

16. SOUTHERN NEW ENGLAND/MID-ATLANTIC YELLOWTAIL FLOUNDER

Larry Alade and Chris Legault

*This assessment of the Southern New England/Mid-Atlantic yellowtail flounder (*Limanda ferruginea*) stock is a Level 2 Management Track assessment of the existing 2012 benchmark assessment (NEFSC 2012). Based on the previous assessment (NEFSC 2022), the stock was overfished, but overfishing was not occurring. This assessment updates commercial fishery catch data, research survey indices of abundance, and the analytical ASAP assessment model and reference points through 2021. Additionally, stock projections have been updated through 2025.*

State of Stock: Based on this updated assessment, the Southern New England/Mid-Atlantic yellowtail flounder (*Limanda ferruginea*) stock is overfished and overfishing is not occurring (Figures 61–62). Retrospective adjustments were made to the model results. Spawning stock biomass (*SSB*) in 2021 was estimated to be 70 mt which is 4% of the biomass target ($SSB_{MSY\ proxy} = 1,715$; Figure 61). The 2021 fully selected fishing mortality was estimated to be 0.082 which is 23% of the overfishing threshold proxy ($F_{MSY\ proxy} = 0.349$; Figure 62).

Table 38: Catch and status table for Southern New England/Mid-Atlantic yellowtail flounder. All weights are in (mt) recruitment is in (000s) and F_{Full} is the fishing mortality on fully selected ages (ages 4 and 5). Model results are from the current updated ASAP assessment. Note: Terminal year estimates of *SSB* and *F* reflect the unadjusted values for retrospective error.

	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
	<i>Data</i>									
Commercial discards	221	185	109	53	26	16	8	6	5	4
Commercial landings	342	461	516	284	126	48	11	2	2	1
Catch for Assessment	563	646	625	337	152	64	19	8	7	5
	<i>Model Results</i>									
Spawning Stock Biomass	1,610	1,318	865	422	159	59	36	45	142	241
F_{Full}	0.585	0.745	0.906	0.961	1.076	1.272	0.788	0.291	0.1	0.032
Recruitment (age-1)	1,671	1,104	179	88	66	179	226	1,155	449	4,396

Projections: Short term projections of biomass were derived by sampling from an empirical cumulative distribution function of 30 recruitment estimates from the ASAP model results. Following the previous and accepted benchmark formulation, recruitment was based on recent estimates of recruitments from the model time series (i.e., corresponding to age-1 in years 1990 through 2019) to reflect the low recent pattern of recruitment in the stock. For projections, the annual fishery selectivity is from the most recent selectivity block in the model, the maturity ogive is the same as assumed for all years in the model,

and mean weights-at-age are from 2014–2019 due to low or no sampling in 2020 and 2021; retrospective adjustments were applied in the projections.

Table 39: Comparison of reference points estimated in an earlier assessment and from the current assessment update. An $F_{40\%SPR}$ proxy was used for the overfishing threshold and was based on long-term stochastic projections.

	2019	2022
$F_{MSY\ proxy}$	0.355	0.349
SSB_{MSY} (mt)	1,756	1,715 (908–2,739)
MSY (mt)	495	461 (245–739)
Median recruitment (age-1) (000s)	6,562	6,004
Overfishing	No	No
Overfished	Yes	Yes

Table 40: Short term projections of total fishery catch and spawning stock biomass for Southern New England/Mid-Atlantic yellowtail flounder based on a harvest scenario of fishing at $F_{MSY\ proxy}$ between 2023 and 2025. Catch in 2022 was assumed to be 4 (mt) based on an estimate provided by the Groundfish Plan Development Team.

Year	Catch (mt)	SSB (mt)	F_{Full}
2022	4	174 (102–293)	0.033 (0.019–0.054)
Year	Catch (mt)	SSB (mt)	F_{Full}
2023	55 (33–91)	203 (121–343)	0.349
2024	84 (46–142)	420 (157–930)	0.349
2025	152 (58–319)	815 (261–1,641)	0.349

Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass, F , recruitment, and population projections).

The long-term outlook for this stock. Recent papers (Stock and Miller 2021, du Pontavice et al. 2022) found a relationship between the Cold Pool Index, a measure of cold water in the Mid-Atlantic Bight, and the stock-recruitment relationship. If the hypothesized relationship holds, and the Cold Pool Index continues to warm due to global climate change, the ability of this stock to support a fishery is questionable. This Management Track assessment followed the approach used in the previous assessment to calculate long-term reference points assuming recruitment remained within the range observed since 1990. If instead recruitment declines in the future, these already low biomass and yield estimates will decrease, providing little potential yield to the fishery.

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted *SSB* or F_{Full} lies outside of the approximate joint confidence region for *SSB* and F_{Full}).

*The 7-year Mohn's ρ , relative to *SSB*, was 0.63 in the 2019 assessment and was 2.43 in 2021. The 7-year Mohn's ρ , relative to *F*, was -0.31 in the 2019 assessment and was -0.62 in 2021. There was a major retrospective pattern for this assessment because the ρ -adjusted estimates of 2021 *SSB* ($SSB_{\rho} = 70$) and 2021 *F* ($F_{\rho} = 0.082$) were outside the approximate 90% confidence regions around *SSB* (148–405) and *F* (0.019–0.053). A retrospective adjustment was made for both the determination of stock status and for projections of catch in 2023. The retrospective adjustment changed the 2021 *SSB* from 241 to 70 and the 2021 F_{Full} from 0.032 to 0.082.*

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?

*Population projections for Southern New England/Mid-Atlantic yellowtail flounder are uncertain for reasons associated with the retrospective pattern and the low stock size. The 2021 estimate of *SSB* is within the bounds of the projected *SSB* from the 2019 assessment, but the ρ -adjusted *SSB* is not. The 2019 estimate of *SSB* from the current assessment is below the 2019 ρ -adjusted *SSB* from the 2019 assessment, indicating the ρ -adjustment applied in the 2019 assessment was not large enough. However, the current low size of this stock makes these comparisons tenuous. This stock is in a rebuilding plan with a rebuilding date of 2029.*

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the effect these changes had on the assessment and stock status.

*Minor changes, in addition to the incorporation of new data, were made to the Southern New England/Mid-Atlantic yellowtail flounder assessment for this update. The larval index was treated differently to now use *SSB* weights-at-age, tuned to the spawning time of the stock instead of the survey time, and used a fixed selectivity pattern equal to the maturity ogive (assumed constant over time in the assessment). These changes more closely reflect the use of the larval index as an indicator of spawning biomass, but had no noticeable impact on the results. The number of fishery selectivity blocks was reduced from 6 to 2 after consideration of a range of alternatives. All of the selectivity block changes produced similar patterns in the residuals (except for using only a single selectivity block), and all showed similar patterns in *F* and *SSB*. The decision to change from 6 to 2 selectivity blocks was based on model parsimony.*

- If the stock status has changed a lot since the previous assessment, explain why this occurred.

The overfishing status of Southern New England/Mid-Atlantic yellowtail flounder has not changed since the last 2019 Management Track assessment. The stock remains at low abundance despite low catches.

- Provide qualitative statements describing the condition of the stock that relate to stock status.

*All three 2021 surveys for Southern New England/Mid-Atlantic yellowtail flounder show record low numbers of fish caught, 2, 3, and 2 in the *NEFSC* spring, *NEFSC* fall, and larval surveys, respectively. The 2022 *NEFSC* spring survey, which is not included in this assessment, caught 3 fish. These surveys were all conducted according to standard operating procedures, so the low numbers of fish caught indicate a low population size, not a problem with the surveys. While low*

fishery catches do not necessarily mean the population is low, the recent catches of < 10 mt in every year since 2019 are consistent with a low population size. There are no indications that this stock is doing well.

- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.

The Southern New England/Mid-Atlantic yellowtail flounder assessment has been used as an example of how to include environmental factors in stock assessments in a number of recent papers (Miller et al. 2016, Xu et al. 2017, Stock and Miller 2021, du Pontavice et al. 2022). All indicate that the environment for this stock is getting worse and causing expected recruitment to decline as the temperature increases in the region. If this trend continues, as expected under nearly all climate models, then the ability of this stock to support a fishery is questionable. Converting the modeling framework for this stock from ASAP to WHAM (or another state-space model) would allow estimation of the relationship between environmental factors and modeled recruitment. The long-term potential yield of this stock associated with climate change could then be considered. A research track assessment for yellowtail flounder stocks in this region is scheduled to begin next month and be peer reviewed in 2024.

- Are there other important issues?

The catchability (q) survey biomass from the Cooperative Research comparative chain sweep experiment (Miller 2013, Jones et al. 2021) estimated similar biomass to the 2021 ρ -adjusted SSB from the assessment, but indicated a much higher biomass than the model-estimated SSB in the 2010s. When these data were used directly in the stock assessment, either the model q was unreasonably high (6 or 10 instead of the expected value of 1) or else there were major problems with the fits to the data. The decline in the expanded survey biomass from 2010 through 2019 cannot easily be explained given the catches and age structure of the fish caught. Future research should explore whether an alternative modeling platform, such as WHAM (Stock and Miller 2021), can find a way to reconcile these data with modeled population estimates.



Limanda ferruginea, Yellowtail Flounder.

16.1. Reviewer Comments: Southern New England/Mid-Atlantic yellowtail flounder

The 2022 assessment for Southern New England/Mid-Atlantic yellowtail flounder (*Limanda ferruginea*) updates the 2012 benchmark assessment use of the ASAP model (NEFSC 2012)²⁶ with additional commercial fishery catch data, five research survey indices of abundance, and reference points through 2019–2021. Stock projections have been updated through 2025.

Retrospective adjustments were made to the ASAP model. Spawning stock biomass in 2021 was estimated to be 70 mt which is 4% of the biomass target ($SSB_{MSY} = 1,715$ mt). The 2021 fully selected fishing mortality was estimated to be 0.082 which is 23% of the overfishing threshold proxy ($F_{MSY proxy} = 0.349$).

Based on this updated assessment, the SNEMA yellowtail flounder stock was overfished but overfishing was not occurring.

Short term projections of biomass were derived by sampling from an empirical cumulative distribution function of 30 recruitment estimates (1990–2019) from the ASAP model results based on recent estimates of recruitments from the model time. The annual fishery selectivity is from the most recent selectivity block in the model, the maturity ogive is the same as assumed for all years in the model, and mean weights-at-age are from 2014–2019 due to low or no sampling in 2020 and 2021. Retrospective adjustments were applied in the projections.

The estimated catch for 2022 is 4 mt, which results in catch advice of 55, 84 and 152 mt for 2023–2025 respectively.

The Panel was very impressed by the amount of work devoted to a stock with its biomass close to the origin.

The Panel was again concerned about the limited sampling for WAA for this stock.

There was concern that the outyear catch advice was overly optimistic given the stock's current biomass and observed failure to recruit. Much of this increase seems to have been fueled by the low fishing pressure and simulated recruitment, which the model responds to by showing sharp increases in biomass. This could be exacerbated by the low sampling in 2020–2021.

Given the status of this stock, NOAA should be sure to provide a timely stock assessment update, and not allow the timing of the species' Research Track effort interfere with this update.

The Panel concluded that the 2022 assessment update for SNEMA yellowtail flounder fulfilled the recommendations of the AOP, is technically sufficient to evaluate stock status and provide scientific advice and meets the Terms of Reference for the stock's assessment. The assessment represents Best Scientific Information Available (BSIA) for this stock for management purposes.

²⁶NEFSC. 2012. 54th Northeast Regional Stock Assessment Workshop (SAW 54) Assessment Report. US Dept Commer, NOAA Fisheries, Northeast Fish Sci Cent Ref Doc. 12-18.; 600p. CRD12-18

References:

- Alade L., 2015. In Northeast Fisheries Science Center. 2015. Operational Assessment of 20 Northeast Groundfish Stocks, Updated Through 2014. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 15-24; 251p. [CRD15-24](#)
- Alade L., 2017. In Northeast Fisheries Science Center. 2017. Operational Assessment of 19 Northeast Groundfish Stocks, Updated Through 2016. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 15-17; 259p. [CRD15-17](#)
- Alade L., C. Legault, S. Cadrin. 2008. In Northeast Fisheries Science Center, 2008. Assessment of 19 Northeast Groundfish Stocks through 2007: Report of the 3rd Groundfish Assessment Review Meeting (**GARM III**), Northeast Fisheries Science Center, Woods Hole, Massachusetts, August 4–8, 2008. US Dept Commer, NOAA Fisheries, Northeast Fish Sci Cent Ref Doc. 08-15; 884p.+xvii. [CRD08-15](#)
- du Pontavice H., T.J. Miller, B.C. Stock, Z. Chen, V.S. Saba. 2022. Ocean model-based covariates improve a marine fish stock assessment when observations are limited. **ICES** Journal of Marine Science. 79: pp.1259–1273. [JMS:fsac050](#)
- Jones A.W., T.J. Miller, P.J. Politis, D.E. Richardson, A.M. Mercer, M.V. Pol, C.D. Roebuck, 2021; Experimental assessment of the effect of net wing spread on relative catch efficiency of four flatfishes by a four seam bottom trawl. Fisheries Research. 244: [106106](#).
- Miller T.J. 2013. A comparison of hierarchical models for relative catch efficiency based on paired-gear data for US northwest Atlantic fish stocks. Canadian Journal of Fisheries and Aquatic Sciences. Vol. 70 (9): pp.1306–1316. [2013-0136](#)
- Miller T.J., J. Hare, L. Alade. 2016. A State-Space approach to Incorporating Environmental Effects on Recruitment in Age-Structured Assessment Model with an Application to Southern New England Yellowtail Flounder. Canadian Journal of Fisheries and Aquatic Sciences. Vol. 73 (8): pp.1261–1270. [2015-0339](#)
- Northeast Fisheries Science Center. 2012. 54th Northeast Regional Stock Assessment Workshop (**SAW 54**) Assessment Report. US Dept Commer, NOAA Fisheries, Northeast Fish Sci Cent Ref Doc. 12-18; 600p. [CRD12-18](#)
- Northeast Fisheries Science Center. 2022. Stock Assessment Update of 14 Northeast Groundfish Stocks Through 2018. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 22-06; 227p. [CRD22-06](#)
- Richardson D., J. Hare, W. Overholtz, D. Johnson. 2009. Development of long-term larval indices for Atlantic herring (*Clupea harengus*) on the northeast US continental shelf. **ICES** Journal of Marine Science, Volume 67, Issue 4, 1 May 2010, pp.617–627. [JMS:fsp276](#)
- Stock B.C., T.J. Miller. 2021. The Woods Hole Assessment Model (**WHAM**): A general state-space assessment framework that incorporates time- and age-varying processes via random effects and links to environmental covariates. Fisheries Research. 240: [105967](#).

Xu H., T.J. Miller, S. Hameed, L.A. Alade, J. Nye. (2017). Evaluating the Utility of the Gulf Stream Index for Predicting Recruitment of Southern New England/Mid-Atlantic yellowtail flounder. *Fisheries Oceanography*. DOI: [10.1111/fog.12236](https://doi.org/10.1111/fog.12236)



Yellowtail Flounder over sandy bottom. Photo credit: iStock




Figure 61: Trends in spawning stock biomass of Southern New England/Mid-Atlantic yellowtail flounder between 1973 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $SSB_{Threshold}$ ($\frac{1}{2}SSB_{MSY proxy}$; horizontal dashed line) as well as SSB_{Target} ($SSB_{MSY proxy}$; horizontal dotted line) based on the 2022 assessment. Biomass was adjusted for a retrospective pattern and the adjustment is shown in red. The approximate 90% log-normal confidence intervals are shown.




Figure 62: Trends in the fully selected fishing mortality (F_{Full}) of Southern New England/Mid-Atlantic yellowtail flounder between 1973 and 2021 from the current (solid line) and previous (dashed line) assessment and the corresponding $F_{Threshold}$ ($F_{MSY proxy} = 0.349$; horizontal dashed line). F_{Full} was adjusted for a retrospective pattern and the adjustment is shown in red; based on the 2022 assessment. The approximate 90% log-normal confidence intervals are shown.




Figure 63: Trends in Recruitment (age-1) (000s) of Southern New England/Mid-Atlantic yellowtail flounder between 1973 and 2021 from the current (solid line) and previous (dashed line) assessment. The approximate 90% log-normal confidence intervals are shown.




Figure 64: Total catch of Southern New England/Mid-Atlantic yellowtail flounder between 1973 and 2021 by fleet (US domestic and foreign catch) and disposition (landings and discards).




Figure 65: Indices of biomass for the Southern New England/Mid-Atlantic yellowtail flounder between 1973 and 2022 for the Northeast Fisheries Science Center (NEFSC) spring and fall bottom trawl surveys. The approximate 90% log-normal confidence intervals are shown. Note: Larval index based on Richardson et al. (2009) was also used in this assessment and is available in the supplemental documentation.

Photo Gallery

Here we provide descriptive text for the photographs and artwork that are scattered throughout the preceding pages.

Archival drawing: American Plaice. Photo [NOAA](#). On page [115](#)

Pollock in a basket. Credit: [NOAA Photo Library](#). On page [128](#)

[NOAA](#) research vessel [Henry B. Bigelow](#), named after Henry Bryant Bigelow (1879–1967), oceanographer and marine biologist. Photo from [NOAA website](#). On page [vii](#)

Sorted Haddock in baskets. Photo [NOAA](#). On page [65](#)

Haddock swimming over rocky bottom. Photo [NOAA](#). On page [53](#)

Atlantic Halibut on deck of fishing vessel. Photo [NOAA](#). On page [75](#)

Monkfish at gravelly sea bottom. Photo credit: iStock. On page [101](#)

Monkfish on the measuring table. Photo: [NOAA Teacher at Sea](#). On page [95](#)

Aerial view of the [NMFS](#) building and surrounds, Woods Hole Laboratory, [MA](#).
The enclosed body of water behind is Eel Pond. Photo [WHOI](#). On page [17](#)

The reason behind it all: seafood display case at a local supermarket. Photo [NOAA](#). On page [iv](#)

Fresh seafood on ice, ready for sale. Credit: Shutterstock. On page [165](#)

Shrimp, mussels, scallop, and fish dish. Credit: iStock. On page [ii](#)

Brian Gay of Millsboro, Delaware holding the record-setting white hake he caught in 2019.
Photo credit: Maryland Department of Natural Resources. On page [86](#)

Aerial view of the buildings and wharves at the Woods Hole Oceanographic Institute, [MA](#).
Two research vessels are docked for re-supply. Photo [WHOI](#). On page [x](#)

Displaying a Winter flounder catch. Photo credit: Willy Goldsmith. On page [42](#)

Wolffish at floor of aquarium tank. Photo credit: Woods Hole Aquarium. On page [21](#)

Yellowtail Flounder over sandy bottom. Photo credit: iStock. On page [158](#)

Hippoglossus hippoglossus, commonly known as Atlantic Halibut, Halibut; range: New England/Mid-Atlantic. Artwork from [NOAA halibut website](#). On page [73](#)

Pollachius virens, commonly known as Atlantic Pollock, Saithe, Coalfish, Coley, Green cod, Boston bluefish; range: New England/Mid-Atlantic. Artwork from [NOAA atlantic pollock website](#). On page 124

Melanogrammus aeglefinus, commonly known as Haddock, Scrod; range: New England/Mid-Atlantic. Artwork from [NOAA haddock website](#). On page 51

Lophius americanus, also known as Goosefish, Monkails, Angler, Fishing frog, Allmouth, Molligut, Abbot, Sea-devil, Lotte; range: New England/Mid-Atlantic, Southeast. Artwork from [NOAA monkfish website](#). On pages 93, 99

Urophycis tenuis, commonly known as White Hake; range: New England/Mid-Atlantic. Artwork from [NOAA white hake website](#). On page 86

Pseudopleuronectes americanus, commonly known as Winter Flounder, Flounder, Sole, Lemon sole, Georges Bank flounder, Blackback flounder; range: New England/Mid-Atlantic, Southeast. Artwork from [NOAA winter flounder website](#). On pages 32, 40

Anarhichas lupus, commonly known as Atlantic wolffish; range: New England/Mid-Atlantic. Artwork from [NOAA wolffish website](#). On page 22

Limanda ferruginea, commonly known as Yellowtail Flounder, Flounder, Rusty dab; range: New England/Mid-Atlantic. Artwork from [NOAA yellowtail flounder website](#). On pages 155



Fresh seafood on ice, ready for sale.

Photo credit: Shutterstock.

Procedures for Issuing Manuscripts in the Northeast Fisheries Science Center Reference Document (CRD) Series and the Technical Memorandum (TM) Series

The mission of NOAA's National Marine Fisheries Service (NMFS) is "stewardship of the nation's ocean resources and their habitat." As the research arm of the NMFS's Greater Atlantic Region, the Northeast Fisheries Science Center (NEFSC) supports the NMFS's mission by "conducting ecosystem-based research and assessments of living marine resources, with a focus on the Northeast Shelf, to promote the recovery and long-term sustainability of these resources and to generate social and economic opportunities and benefits from their use." Results of NEFSC research are largely reported in primary scientific media (e.g., anonymously-peer-reviewed scientific journals). However, to assist itself in providing data, information, and advice to its constituents, the NEFSC occasionally releases its results in its own media.

NOAA Technical Memorandum NMFS-NE—This series is issued irregularly. The series typically includes: data reports of long-term field or lab studies of important species or habitats; synthesis reports for important species or habitats; annual reports of overall assessment or monitoring programs; manuals describing program-wide surveying or experimental techniques; literature surveys of important species or habitat topics; proceedings and collected papers of scientific meetings; and indexed and/or annotated bibliographies. All issues receive internal scientific review and most issues receive technical and copy editing.

NORTHEAST FISHERIES SCIENCE CENTER REFERENCE DOCUMENT—This series is issued irregularly. The series typically includes: data reports on field and lab studies; progress reports on experiments, monitoring, and assessments; background papers for, collected abstracts of, and/or summary reports of scientific meetings; and simple bibliographies. Issues receive internal scientific review and most issues receive copy editing.

Clearance

All manuscripts submitted for issuance as **CRD**s must have cleared the **NEFSC**'s manuscript/abstract/webpage review process. If your manuscript includes material from another work which has been copyrighted, then you will need to work with the **NEFSC**'s Editorial Office to arrange for permission to use that material by securing release signatures on the "NEFSC Use-of-Copyrighted-Work Permission Form."

For more information, **NEFSC** authors should see the **NEFSC**'s online publication policy manual, "Manuscript/abstract/webpage preparation, review, and dissemination: **NEFSC** Author's Guide to Policy, Process, and Procedure."

Style

The **CRD** series is obligated to conform with the style contained in the current edition of the United States Government Printing Office Style Manual; however that style manual is silent on many aspects of scientific manuscripts. The **CRD** series relies more on the **CSE** Style Manual. Manuscripts should be prepared to conform with both of these style manuals.

The **CRD** series uses the Integrated Taxonomic Information System, the American Fisheries Society's guides and the Society for Marine Mammalogy's guide for verifying scientific species names.

For in-text citation, use the name-date system. A special effort should be made to ensure that all necessary bibliographic information is included in the list of references cited. Personal communications must include date, full name, and full mailing address of the contact.

Preparation

Once your document has cleared the review process, the Editorial Office will contact you with publication needs — for example, revised text (if necessary) and separate digital figures and tables if they are embedded in the document. Materials may be submitted to the Editorial Office as email attachments or intranet downloads. Text files should be in Microsoft Word, tables may be in Word or Excel, and graphics files may be in a variety of formats (JPG, GIF, Excel, PowerPoint, etc.).

Production and Distribution

The Editorial Office will perform a copy-edit of the document and may request further revisions. The Editorial Office will develop the inside and outside front covers, the inside and outside back covers, and the title and bibliographic control pages of the document.

Once the **CRD** is ready, the Editorial Office will contact you to review it and submit corrections or changes before the document is posted online. A number of organizations and individuals in the Northeast Region will be notified by e-mail of the availability of the document online.
